For Research Use Only

GAD1 Polyclonal antibody

Catalog Number:10408-1-AP

Featured Product

24 Publications




Basic Information	Catalog Number: 10408-1-AP	GenBank Accession Number: BC002815	Purification Method: Antigen affinity purification	
	Size:	GeneID (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 600 µg/ml by		WB 1:5000-1:50000	
	Nanodrop;	UNIPROT ID:	IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	Source:	Q99259	protein lysate IHC 1:200-1:800	
	Rabbit	Full Name:	IIIC 1.200-1.000	
	Isotype: IgG	glutamate decarboxylase 1 (brain, 67kDa)		
	Immunogen Catalog Number: AG0666	Calculated MW: 67 kDa		
		Observed MW: 67 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IP, IHC, ELISA	WB : mouse brain tissue, rat brain tissue		
	Cited Applications: WB, IF, IHC	IP : mouse b	IP : mouse brain tissue,	
	Species Specificity: human, mouse, rat	IHC : mouse	IHC : mouse brain tissue,	
	Cited Species: human, rat, mouse			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
	The GAD1(glutamate decarboxylase 1) gene encodes 67 kDa glutamic acid decarboxylase isoform (GAD67), the rate-limiting enzyme responsible for γ-aminobutyric acid (GABA) biosynthesis from glutamatic acid and the majo GAD isoform in the human brain for early brain development (PMID:21302352). It is also potentially involved in variety of skin activities. It belongs to the group II decarboxylase family.			
Background Information	rate-limiting enzyme responsible fo GAD isoform in the human brain for	r γ -aminobutyric acid (GABA) biosynt early brain development (PMID:2130)	hesis from glutamatic acid and the maj	
	rate-limiting enzyme responsible fo GAD isoform in the human brain for o variety of skin activities. It belongs t	r γ-aminobutyric acid (GABA) biosyn early brain development (PMID:2130 to the group II decarboxylase family.	hesis from glutamatic acid and the maj 2352). It is also potentially involved in	
	rate-limiting enzyme responsible for GAD isoform in the human brain for evariety of skin activities. It belongs to Author Pu	r γ-aminobutyric acid (GABA) biosym early brain development (PMID:2130) to the group II decarboxylase family. bmed ID Journal	thesis from glutamatic acid and the major 2352). It is also potentially involved in Application	
Background Information	rate-limiting enzyme responsible for GAD isoform in the human brain for variety of skin activities. It belongs to Author Pu Yifei Xu 36	r γ-aminobutyric acid (GABA) biosyn early brain development (PMID:2130) to the group II decarboxylase family.	hesis from glutamatic acid and the maj 2352). It is also potentially involved in	
	rate-limiting enzyme responsible for GAD isoform in the human brain for variety of skin activities. It belongs to Author Pu Yifei Xu 36 Lei Wu 33	r γ -aminobutyric acid (GABA) biosyn early brain development (PMID:2130) to the group II decarboxylase family. bmed ID Journal 120289 Front Pharmacol	hesis from glutamatic acid and the maj 2352). It is also potentially involved in Application WB	
	rate-limiting enzyme responsible for GAD isoform in the human brain for variety of skin activities. It belongs to Author Pu Yifei Xu 36 Lei Wu 33	r y -aminobutyric acid (GABA) biosyn early brain development (PMID:2130) to the group II decarboxylase family. bmed ID Journal 120289 Front Pharmacol 044778 Phytother Res 467583 Transl Psychiatry ter shipment.	hesis from glutamatic acid and the maj 2352). It is also potentially involved in Application WB	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10408-1-AP (GAD1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



IP result of anti-GAD1 (IP:10408-1-AP, 4ug; Detection:10408-1-AP 1:500) with mouse brain tissue lysate 1920 ug.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 10408-1-AP (GAD1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 10408-1-AP (GAD1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).