For Research Use Only

GORASP2 Polyclonal antibody

Catalog Number:10598-1-AP

Featured Product

60 Publications




Size: 150ul , Concentration: 450 µg/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number:	GeneID (NCBI): 26003 UNIPROT ID: Q9H8Y8 Full Name: golgi reassembly s 55kDa	to duing protoin 2	Recommended Dilutions: WB 1:500-1:3000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:20-1:200	
AG0950	Calculated MW: 55 kDa Observed MW: 59 kDa	tacking protein 2,	IF 1:50-1:500	
Tested Applications: WB, IP, IF, IHC, ELISA Cited Applications:			rols: O cells, rat testis tissue, HeLa cells, Raj	
WB, IF, IHC				
Species Specificity:		IHC : human colon cancer tissue,		
	IF : HenG2 co		ells,	
human, mouse, monkey				
TE buffer pH 9.0; (*) Alternati	ively, antigen			
phosphorylated by MKK/ERK pathwa stacking of the Golgi cisternae, and	y during mitoisis. GC in the process by whi	RASP2 plays a rol ch Golgi stacks ref	e in the assembly and membrane orm after mitotic breakdown.GORASP	
Author Pu	bmed ID 10	ırnal	Application	
			IF	
Xihua Yue 34	493279 BM	C Biol	WB	
Myun Hwa Dunlop 28	874656 Na	t Commun		
Storage Buffer: PBS with 0.02% sodium azide and 50	0% glycerol pH 7.3.			
	WB, IP, IF, IHC, ELISA Cited Applications: WB, IF, IHC Species Specificity: human, mouse, rat Cited Species: human, mouse, monkey Note-IHC: suggested antigen TE buffer pH 9.0; (*) Alternative retrieval may be performed ve buffer pH 6.0 GORASP2 (Golgi reassembly-stacking phosphorylated by MKK/ERK pathwas stacking of the Golgi cisternae, and i supposed to regulate the intracellulity as transmembrane TGFA. Author Pu Jennifer Y Liu 34 Xihua Yue 34 Myun Hwa Dunlop 28 Storage: Storage Storage Store at -20°C. Stable for one year af Storage Buffer: PBS with 0.02% sodium azide and 50	Tested Applications: WB, IP, IF, IHC, ELISA Cited Applications: WB, IF, IHC Species Specificity: human, mouse, rat Cited Species: human, mouse, monkey Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 GORASP2 (Golgi reassembly-stacking protein 2, or GRS2) phosphorylated by MKK/ERK pathway during mitoisis. GO stacking of the Golgi cisternae, and in the process by whi supposed to regulate the intracellular transport and prese as transmembrane TGFA. Author Pubmed ID Jou Jennifer Y Liu 34533192 Bic Xihua Yue Myun Hwa Dunlop 28874656 Na Storage: Storage: Storage at -20°C. Stable for one year after shipment.	Tested Applications:Positive ContinuesWB, IP, IF, IHC, ELISAWB : COLO 32Cited Applications:cellsWB, IF, IHCIP : HeLa cellsSpecies Specificity:IHC : human chuman, mouse, ratIF : HepG2 cellhuman, mouse, monkeyIF : HepG2 cellNote-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0GORASP2 (Golgi reassembly-stacking protein 2, or GRS2) has multiple phos phosphorylated by MKK/ERK pathway during mitoisis. GORASP2 plays a role stacking of the Golgi cisternae, and in the process by which Golgi stacks refe supposed to regulate the intracellular transport and presentation of a define as transmembrane TGFA.AuthorPubmed IDJournalJennifer Y Liu34493279BMC BiolXihua Yue34493279BMC BiolMyun Hwa Dunlop28874656Nat CommunStorage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3.	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10598-1-AP (GORASP2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




WB result of GORASP2 antibody (10598-1-AP; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-GORASP2 transfected HeLa cells.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using GORASP2 antibody (10598-1-AP) at dilution of 1:200 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL555-Phalloidin (orange).



Immunohistochemical analysis of paraffinembedded human colon cancer using 10598-1-AP (GORASP2 antibody) at dilution of 1:100 (under 10x lens).



IP result of anti-GORASP2 (IP:10598-1-AP, 3ug; Detection:10598-1-AP 1:700) with HeLa cells lysate 1320ug.