For Research Use Only

NRAS Polyclonal antibody

Catalog Number:10724-1-AP

Featured Product






Basic Information	Catalog Number: 10724-1-AP	GenBank Accession Number: BC005219	Purification Method: Antigen affinity purification		
	Size:	GeneID (NCBI):	Recommended Dilutions:		
	150ul, Concentration: 273 µg/ml by Bradford method using BSA as the standard; Source:	4893	WB 1:500-1:2000		
		UNIPROT ID:	IHC 1:50-1:500 IF 1:50-1:500		
		P01111	11 1.90 1.900		
	Rabbit Isotype: IgG Immunogen Catalog Number: AG1081	Full Name: neuroblastoma RAS viral (v-ras) oncogene homolog Calculated MW:			
				21 kDa	
				Observed MW:	
		21 kDa			
		Applications	Tested Applications:	Positive Controls:	
			WB, IF, FC, IHC, ELISA	WB : A43	WB : A431 cells, C6 cells, NIH/3T3 cells, Jurkat cells, A549 cells, HEK-293 cells, MCF-7 cells
Cited Applications:	A 549 cel				
			nan pancreas tissue, human testis tissue		
Species Specificity: human, mouse, rat	IF : HeLa cells,				
Cited Species:					
human, mouse					
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
Background Information	ras gene family and it is a member of transduction pathways, and bind GDI	of the small GTPase superfamily. R P/GTP and possess intrinsic GTPas	AS proteins are involved in signal		
	ras gene family and it is a member of transduction pathways, and bind GDI it is activated in HL60, a promyelocy leukemia (JMML).	of the small GTPase superfamily. R P/GTP and possess intrinsic GTPas	e activity. It is mapped on chromosome 1, a		
Background Information	ras gene family and it is a member of transduction pathways, and bind GDI it is activated in HL60, a promyelocy leukemia (JMML).	f the small GTPase superfamily. R P/GTP and possess intrinsic GTPas tic leukemia line. Defects in NRAS	AS proteins are involved in signal e activity. It is mapped on chromosome 1, a are a cause of juvenile myelomonocytic		
	ras gene family and it is a member of transduction pathways, and bind GDI it is activated in HL60, a promyelocy leukemia (JMML). Author Pu Andrew M Waters 28	of the small GTPase superfamily. R P/GTP and possess intrinsic GTPas tic leukemia line. Defects in NRAS bmed ID Journal	AS proteins are involved in signal e activity. It is mapped on chromosome 1, a s are a cause of juvenile myelomonocytic Application		
	ras gene family and it is a member of transduction pathways, and bind GDI it is activated in HL60, a promyelocy leukemia (JMML). Author Pu Andrew M Waters 28 Akiya Tatsumi 36	of the small GTPase superfamily. R P/GTP and possess intrinsic GTPas tic leukemia line. Defects in NRAS bmed ID Journal 951536 Sci Signal	AS proteins are involved in signal e activity. It is mapped on chromosome 1, a s are a cause of juvenile myelomonocytic Application WB		
	ras gene family and it is a member of transduction pathways, and bind GDI it is activated in HL60, a promyelocy leukemia (JMML). Author Pu Andrew M Waters 28 Akiya Tatsumi 36	of the small GTPase superfamily. R P/GTP and possess intrinsic GTPas tic leukemia line. Defects in NRAS bmed ID Journal 951536 Sci Signal 358254 Biology (Basel) 046519 Clin Cancer Res ter shipment.	AS proteins are involved in signal e activity. It is mapped on chromosome 1, a are a cause of juvenile myelomonocytic Application WB WB		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10724-1-AP (NRAS antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human pancreas tissue slide using 10724-1-AP (NRAS Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human pancreas tissue slide using 10724-1-AP (NRAS Antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using 10724-1-AP (NRAS antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human NRAS (10724-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP, Clone:) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).