For Research Use Only

Bcl-XL Polyclonal antibody

Catalog Number: 10783-1-AP

Featured Product

172 Publications



Basic Information

Catalog Number:

10783-1-AP

Size:

GenBank Accession Number:

BC019307

GeneID (NCBI):

150ul, Concentration: 450 µg/ml by 598

Nanodrop; UNIPROT ID:

Source: Q07817
Rabbit Full Name:
Isotype: BCL2-like 1
IgG Calculated MW:

Immunogen Catalog Number: 26 kDa AG1232

.232 Observed MW:

30 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:2000-1:12000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF 1:10-1:100

Applications

Tested Applications:

WB, IP, IF, FC, IHC, ELISA

Cited Applications:

WB, IP, IF, IHC, ChIP

Species Specificity: human, mouse, rat

Cited Species:

human, goat, rat, sheep, mouse, zebrafish, hamster, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A549 cells, Raji cells, Jurkat cells, rat brain tissue, HeLa cells, NIH/3T3 cells, human placenta tissue, C6 cells, K-562 cells, mouse placenta tissue, RAW264.7

IP: k-562 cells,

IHC: human lung cancer tissue, human colon tissue

IF: NIH/3T3 cells,

Background Information

BCL2L1 is a member of the BCL-2 protein family. BCL2L1 is expressed as three isoforms, Bcl-X(L), Bcl-X(s), and Bcl-X(beta), and is located at the outer mitochondrial membrane. The Bcl-X(L) isoform is a 233 amino acid protein, acting as an apoptotic inhibitor. Bcl-XL can form heterodimers with BAX, BAK, or BCL2, and the heterodimerization with BAX does not seem to be required for anti-apoptotic activity. The Bcl-X(s) isoform is a shorter variant that is 178 amino acids in length and lacks a 63 amino acid region (amino acids 126-188), acting as an apoptotic activator. Bcl-X(beta) is a 227 amino acid protein. This antibody can recognize Bcl-XL, Bcl-X(s) and Bcl-X(Beta).

Notable Publications

Author	Pubmed ID	Journal	Application
Yanhui Zhu	36183949	J Ethnopharmacol	IF
Xufeng Tao	25083618	Transplantation	WB
Yanling Lin	36139911	Antioxidants (Basel)	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10783-1-AP (Bcl-XL antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




WB result of Bcl-XL antibody (10783-1-AP, 1:800) with si-Control and si-Bcl-XL transfected HeLa cells.




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 10783-1-AP (Bcl-XL antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 10783-1-AP (Bcl-XL antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of () fixed NIH/3T3 cells using 10783-1-AP (Bcl-XL antibody) at dilution of 1:25 and Rhodamine-Goat anti-Rabbit IgG.



IP result of anti-Bcl-XL (IP:10783-1-AP, 4ug; Detection:10783-1-AP 1:1000) with k-562 cells lysate 1760 ug.



1X10^6 Jurkat cells were intracellularly stained with 0.2 ug Anti-Human Bcl-XL (10783-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (green), and 0.2 ug Control Antibody. Cells were fixed with 90% MeOH.