

For Research Use Only

P16-INK4A Polyclonal antibody

Catalog Number: 10883-1-AP

Featured Product

319 Publications



Basic Information

Catalog Number:

10883-1-AP

Size:

150ul, Concentration: 650 µg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1328

GenBank Accession Number:

BC021998

GeneID (NCBI):

1029

UNIPROT ID:

Q8N726

Full Name:

cyclin-dependent kinase inhibitor 2A (melanoma, p16, inhibits CDK4)

Calculated MW:

16 kDa

Observed MW:

16-18 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:8000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:1000-1:4000

Applications

Tested Applications:

WB, IP, FC, ELISA

Cited Applications:

WB, IP, FC, Cell treatment

Species Specificity:

human

Cited Species:

human, monkey, canine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HEK-293 cells, HEK293 cells, HeLa cells, HepG2 cells, PC-3 cells

IP : HEK-293 cells,

IHC : human cervical cancer tissue,

Background Information

CDKN2A generates several transcript variants which differ in their first exons. At least three alternatively-spliced variants encoding distinct proteins were reported. Two of them named p16-INK4 and p14 are sharing 50% identity. The third one -p14(ARF) is entirely unrelated. 10883-1-AP reacts with p16 specifically. P16 plays an essential role in regulating the cell cycle, and mutations in p16 increase the risk of developing various cancers, including melanoma.

Notable Publications

Author	Pubmed ID	Journal	Application
Shin Hamada	28971839	Am J Physiol Gastrointest Liver Physiol	
Julie Wang	26416809	Circulation	WB
Shengya Tian	31562192	Life Sci Alliance	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10883-1-AP (P16-INK4A antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




WB result of p16-INK4A antibody (10883-1-AP, 1:2,000) with si-Control and si-p16 transfected HEK-293 cells.




Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 10883-1-AP (P16-INK4A antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-P16-INK4A (IP:10883-1-AP, 4ug; Detection:10883-1-AP 1:4000) with HEK-293 cells lysate 1400 ug.



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 10883-1-AP (P16-INK4A antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10⁶ HeLa cells were intracellularly stained with 0.5 ug Anti-Human P16-INK4A (10883-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.5 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).