## For Research Use Only

## BID Polyclonal antibody

Catalog Number: 10988-1-AP

**Featured Product** 

54 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

10988-1-AP BC009197 GeneID (NCBI): Size:

150ul , Concentration: 700 µg/ml by Nanodrop and 227  $\mu$ g/ml by Bradford UNIPROT ID: method using BSA as the standard; P55957

Source: Full Name:

Rabbit BH3 interacting domain death agonist

Isotype: Calculated MW: 22 kDa Immunogen Catalog Number: Observed MW: AG1445 22 kDa

**Applications** 

**Tested Applications:** 

WB, IP, ELISA Cited Applications:

WB, IHC

Species Specificity:

human **Cited Species:** human, rat, mouse, pig Positive Controls:

WB: A431 cells, human brain tissue, Tunicamycin treated HeLa cells, A549 cells, HEK-293 cells, HepG2 cells, Jurkat cells

**Purification Method:** 

WB 1:1000-1:4000

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

IP: HeLa cells,

**Background Information** 

BID, also named as p22 BID, can be cleaved into p11 BID, p13 BID and p15 BID. It is pro-apoptotic molecules. The major proteolytic product p15 BID allows the release of cytochrome c. BID-L, BID-EL and BID-ES induce ICE-like proteases and apoptosis. BID-S does not induce apoptosis. BID counters the protective effect of Bcl-2. (PMID:14583606)

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Bin Xu	31637186	Int J Ophthalmol	WB
Tahir Mehmood	29151958	J Cancer	WB
Robert Pedley	33067418	Cell Death Dis	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10988-1-AP (BID antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



WB result of BID antibody (10988-1-AP, 1:8000) with si-Control and si-BID transfected HeLa cells.



Tunicamycin treated HeLa cells were subjected to SDS PAGE followed by western blot with 10988-1-AP (BID Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-BID (IP:10988-1-AP, 4ug; Detection:10988-1-AP 1:500) with HeLa cells lysate 1280 ug.