

For Research Use Only

p63 Polyclonal antibody

Catalog Number: 12143-1-AP

Featured Product

29 Publications



Basic Information

Catalog Number:

12143-1-AP

Size:

150ul, Concentration: 550 µg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG2791

GenBank Accession Number:

BC039815

GeneID (NCBI):

8626

UNIPROT ID:

Q9H3D4

Full Name:

tumor protein p63

Calculated MW:

680 aa, 77 kDa

Observed MW:

63-68 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:200-1:800

IF 1:400-1:1600

Applications

Tested Applications:

WB, IP, IF, FC, ELISA

Cited Applications:

WB, IP, IF, ColP, ChIP

Species Specificity:

human, mouse, rat

Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A431 cells, human heart tissue, mouse heart tissue, U-937 cells, Apoptosised HeLa cells, HaCaT cells, PC-12 cells, mouse thymus tissue

IP: A431 cells,

IHC: human tonsillitis tissue, human prostate cancer tissue

IF: A431 cells, human lung cancer tissue

Background Information

TP63, also named KET, P63, P73H, P73L and TP73L, belongs to the p53 family. It is a homologue of the tumor suppressor p53 and p73 genes. It is involved in malignancy acquisition and maintenance of cells. Unlike p53, the p63 gene encodes multiple isoforms with remarkably divergent abilities to transactivate p53 reporter genes and induce apoptosis. TP63 acts as a sequence specific DNA binding transcriptional activator or repressor. TP63 has 12 isoforms with MW 40kd(P40), 50kd(P60), 63kd(P63) and 73kd(P73L). It is Nuclear stain.

Notable Publications

Author	Pubmed ID	Journal	Application
Maximilian Haas	31553905	Cell Rep	IF
Vincent Manna	34585152	STAR Protoc	IF
Xiafang Wu	29081891	Oxid Med Cell Longev	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12143-1-AP (p63 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 12143-1-AP (p63 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 12143-1-AP (p63 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using p63 antibody (12143-1-AP) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



IP result of anti-p63 (IP:12143-1-AP, 4ug; Detection:12143-1-AP 1:1000) with A431 cells lysate 1200 ug.



1×10^6 A431 cells were intracellularly stained with 0.4 ug Anti-Human p63 (12143-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).