For Research Use Only

PMP2 Polyclonal antibody

Catalog Number: 12717-1-AP

17 Publications



Basic Information

Catalog Number: GenBank Accession Number: 12717-1-AP BC034997

GeneID (NCBI): Size

150ul, Concentration: 1000 µg/ml by 5375 Nanodrop and 540 µg/ml by Bradford Full Name:

method using BSA as the standard; peripheral myelin protein 2

Calculated MW: Rabbit 132 aa, 15 kDa Isotype: Observed MW: IgG 15-20 kDa

Immunogen Catalog Number:

AG3411

Applications

Tested Applications: IHC, WB,ELISA

Cited Applications: IF, IHC, WB

Species Specificity: human, mouse, rat, pig

Cited Species: human, rat, mouse, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: pig spinal cord tissue, human brain tissue, mouse

Purification Method:

WB 1:500-1:2000

IHC 1:20-1:200

Antigen affinity purification

Recommended Dilutions:

testis tissue

IHC: human gliomas tissue,

Background Information

PMP2 (peripheral myelin protein-2, also known as P2) is one of the major proteins of peripheral myelin and appears to be related to the transport of fatty acids or the metabolism of myelin lipids.P2 constitutes up to 15% of total protein of peripheral nervous system (PNS) myelin. It is also present in small amounts in central nervous system (CNS) myelin, being abundant in spinal cord and brain stem myelin. As a structural protein, P2 is thought to stabilize the myelin membranes.

Notable Publications

Author	Pubmed ID	Journal	Application
Lakshmi Dhevi N Selvan	26181685	Mol Biosyst	IHC
Jureepon Roboon	36274991	Front Cell Neurosci	WB
Jennifer Zenker	24849898	Glia	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


*** 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human gliomas using 12717-1-AP (PMP2 antibody) at dilution of 1:100 (under 10x lens).



pig spinal cord tissue were subjected to SDS PAGE followed by western blot with 12717-1-AP (PMP2 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.