

For Research Use Only

TDP-43 (C-terminal) Polyclonal antibody



Catalog Number: 12892-1-AP

Featured Product

220 Publications

Basic Information

Catalog Number:

12892-1-AP

Size:

150ul, Concentration: 650 µg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC001487

GeneID (NCBI):

23435

UNIPROT ID:

Q13148

Full Name:

TAR DNA binding protein

Calculated MW:

43 kDa

Observed MW:

43-45 kDa, 35 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:5000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:500-1:2000

IF 1:2000-1:8000

Applications

Tested Applications:

WB, IP, IF, IHC, ELISA

Cited Applications:

WB, IP, IF, IHC, CoIP, ChIP

Species Specificity:

human, mouse, rat

Cited Species:

human, chicken, rat, mouse, monkey, zebrafish, Drosophila

Positive Controls:

WB: A549 cells, mouse brain tissue, HeLa cells, C6 cells

IP: HeLa cells, mouse brain tissue

IHC: mouse brain tissue, human gliomas tissue

IF: HeLa cells, mouse brain tissue, Neuro-2a cells

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Transactivation response (TAR), DNA-binding protein of 43 kDa (also known as TARDBP or TDP-43), was first isolated as a transcriptional inactivator binding to the TAR DNA element of the HIV-1 virus. Neumann et al. (2006) found that a hyperphosphorylated, ubiquitinated, and cleaved form of TARDBP, known as pathologic TDP-43, is the major component of the tau-negative and ubiquitin-positive inclusions that characterize amyotrophic lateral sclerosis (ALS) and the most common pathological subtype of frontotemporal lobar degeneration (FTLD-U). 12892-1-AP is a rabbit polyclonal antibody raised against the C-terminal amino acids of human TDP-43. This antibody recognizes the cleavage product of 20-30 kDa in addition to the native and phosphorylated forms of TDP-43. Immunohistochemical analyses of TDP-43 using this antibody detect both normal diffuse nuclear staining and insoluble inclusions in pathologic tissues. Various forms of TDP-43 exist, including 18-35 kDa of cleaved C-terminal fragments, 45-50 kDa phosphoprotein, 55 kDa glycosylated form, 75 kDa hyperphosphorylated form, and 90-300 kDa cross-linked form. (17023659,19823856,21666678,22193176)

Recently TDP-43 has been reported to be overexpressed in triple negative breast cancer (TNBC) and it may be a potential target for TNBC diagnosis and drug design. (29581274)

Notable Publications

Author	Pubmed ID	Journal	Application
Helena Gossye	36171642	Brain	IHC
Y Liu	25270903	Neuroscience	
Wenrui Huang	34580300	Nat Commun	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 12892-1-AP (TDP-43 (C-terminal) antibody at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using TDP-43 (C-terminal) antibody (12892-1-AP) at dilution of 1:4000 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).




A549 cells (shcontrol and shRNA of TDP43) were subjected to SDS PAGE followed by western blot with 12892-1-AP (TDP43 antibody) at dilution of 1:1000.



HeLa cells were subjected to SDS PAGE followed by western blot with 12892-1-AP (TDP-43 (C-terminal) antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



IP result of anti-TDP-43 (C-terminal) (IP:12892-1-AP, 3ug; Detection:12892-1-AP 1:1000) with HeLa cells lysate 3000ug.



Immunofluorescent analysis of unfixed frozen OCT-embedded mouse brain tissue using TDP-43 (C-terminal) antibody (12892-1-AP) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).