### For Research Use Only

# NF-L Polyclonal antibody

Catalog Number: 12998-1-AP

**20 Publications** 



### **Basic Information**

Catalog Number: GenBank Accession Number: 12998-1-AP BC039237

ze: GenelD (NCBI):

150ul , Concentration: 600 µg/ml by 4747 Nanodrop; Full Name

Source: neurofilament, light polypeptide

Rabbit Calculated MW:

Isotype: 543 aa, 62 kDa

IgG Observed MW:

Immunogen Catalog Number: 65-68 kDa, 140-160 kDa

AG3667

# **Applications**

**Tested Applications:** 

FC, IF, IHC, IP, WB, ELISA
Cited Applications:

IF, IHC, IP, WB Species Specificity: human, mouse, rat

Cited Species: human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

## Purification Method:

Antigen affinity purification

#### Recommended Dilutions:

WB 1:2000-1:16000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:1000-1:4000 IF 1:200-1:800

#### Positive Controls:

WB: SH-SY5Y cells, Neuro-2a cells, mouse brain tissue,

C6 cells

IP: mouse brain tissue,

IF: SH-SY5Y cells, mouse brain tissue

# **Background Information**

NEFL, also named as NF68 and NF-L, belongs to the intermediate filament family. Neurofilaments are the 10 nm intermediate filaments found specifically in neurons. They are a major component of the cell's cytoskeleton, and provide support for normal axonal radial growth. Neurofilaments usually contain three intermediate filament proteins: L, M, and H, which are involved in the maintenance of neuronal caliber. The names given to the three major neurofilament subunits are based upon the apparent molecular weight of the mammalian subunits on SDS-PAGE: NF-L, 65-68 kDa; NF-M, 145-160 kDa and NF-H, 200-220 kDa. This antibody is specific to NEFL It has a very weak reaction to NEFM.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Xiaoyin Liu	36246376	Front Bioeng Biotechnol	IF
Brendan P Major	33117257	Front Neurol	
Fei Yin	25374587	Neural Regen Res	WB

# Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


### Selected Validation Data




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 12998-1-AP (NF-L antibody) at dilution of 1:2000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 12998-1-AP (NF-L antibody) at dilution of 1:2000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




WB result of 12998-1-AP.




IP Result of anti-NF-L (IP:12998-1-AP, 4ug; Detection:12998-1-AP 1:1000) with mouse brain tissue lysate 4000ug.




1x10^6 SH-SY5Y cells were intracellularly stained with 0.4 ug Anti-Human NF-L (12998-1-AP) and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).




Various lysates were subjected to SDS PAGE followed by western blot with 12998-1-AP (NF-L antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using NF-L antibody (12998-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using NF-L antibody (12998-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using NF-L antibody (12998-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).