

For Research Use Only

p115, USO1 Polyclonal antibody

Catalog Number: 13509-1-AP

Featured Product

19 Publications



Basic Information

Catalog Number: 13509-1-AP	GenBank Accession Number: BC032654	Purification Method: Antigen affinity purification
Size: 150ul, Concentration: 550 µg/ml by Nanodrop;	GeneID (NCBI): 8615	Recommended Dilutions: WB 1:1000-1:8000 IP 0.5-4.0 ug for IP and 1:500-1:3000 for WB
Source: Rabbit	Full Name: USO1 homolog, vesicle docking protein (yeast)	IHC 1:50-1:500 IF 1:500-1:2000
Isotype: IgG	Calculated MW: 962 aa, 108 kDa	
Immunogen Catalog Number: AG4431	Observed MW: 115 kDa	

Applications

Tested Applications:
IF, IHC, IP, WB, ELISA

Cited Applications:
IF, WB

Species Specificity:
human, mouse, rat

Cited Species:
human, mouse

Positive Controls:

WB: HEK-293 cells, mouse testis tissue, mouse thymus tissue, HeLa cells, human brain tissue, SH-SY5Y cells, HepG2 cells

IP: mouse brain tissue,

IHC: human gliomas tissue,

IF: HeLa cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

p115, also known as USO1, TAP (transcytosis-associated protein) or VDP (vesicle docking protein) is a general vesicular transport factor and plays an important role at different steps of vesicular transport. It is a 962-residue peripheral membrane protein which recycles between the cytosol and the Golgi apparatus during interphase (PMID: 9478999). p115 forms stable homodimers (PMID: 19247479). Rab1 recruits p115 to coat protein complex II (COPII) vesicles during budding from the endoplasmic reticulum, where p115 interacts directly with a select set of SNARE proteins (PMID: 10903204). p115 is required for intra-Golgi transport, and also functions in endoplasmic reticulum to Golgi trafficking, Golgi biogenesis and exocytotic transport (PMID: 19247479).

Notable Publications

Author	Pubmed ID	Journal	Application
Mohsan Saeed	32997711	PLoS Pathog	WB
Guillermo Arango Duque	34580108	J Immunol	IF
Jing Wang	29025970	J Cell Sci	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of p115, USO1 antibody (13509-1-AP; 1:40000); incubated at room temperature for 1.5 hours) with sh-Control and sh-p115, USO1 transfected HeLa cells.




Various lysates were subjected to SDS PAGE followed by western blot with 13509-1-AP (p115, USO1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using p115, USO1 antibody (13509-1-AP) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 13509-1-AP (p115, USO1 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 13509-1-AP (p115, USO1 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP Result of anti-p115, USO1 (IP:13509-1-AP, 3ug; Detection:13509-1-AP 1:1500) with mouse brain tissue lysate 7000ug.