For Research Use Only

CD80/B7-1 Polyclonal antibody

Catalog Number:14292-1-AP

25 Publications



Basic Information

Catalog Number:

14292-1-AP

Nanodrop:

GenBank Accession Number:

BC042665

GeneID (NCBI):

150ul , Concentration: 700 $\mu g/ml$ by

UNIPROT ID:

P33681 Rabbit Isotype:

Full Name: CD80 molecule

IgG Immunogen Catalog Number: Calculated MW: 33 kDa

AG5615

Observed MW:

60 kDa

Applications

Tested Applications:

WB, FC, ELISA

Cited Applications:

WB. FC

Species Specificity: human, mouse

Cited Species: human, mouse **Positive Controls:**

WB: Raji cells, RAW 264.7 cells, human peripheral

Purification Method:

WB 1:500-1:2000

Antigen affinity purification

Recommended Dilutions:

blood leukocyte, Ramos cells

Background Information

CD80 (also known as B7-1) is a type I membrane protein that is a member of the immunoglobulin superfamily, with an extracellular immunoglobulin constant-like domain and a variable-like domain required for receptor binding. It is expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, monocytes, and macrophages. CD80 is the receptor for the proteins CD28 and CTLA-4 found on the surface of T-cells. It is involved in the $costimulatory\ signal\ essential\ for\ T-lymphocyte\ activation.\ T-cell\ proliferation\ and\ cytokine\ production\ is\ induced$ by the binding of CD28, binding to CTLA-4 has opposite effects and inhibits T-cell activation. CD80 also acts as a cellular attachment receptor for adenovirus subgroup B. (PMID: 7545666; 12015893; 16920215)

Notable Publications

Author	Pubmed ID	Journal	Application
Haoran Wang	35087966	Bioact Mater	FC
Wen Song	36260348	J Med Chem	FC
Jiayu Zhang	35530972	Int J Nanomedicine	

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 14292-1-AP (CD80/B7-1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



1X10^6 Raji cells were stained with 0.2ug CD80/B7-1 antibody (14292-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.