

For Research Use Only

NFKB2,p52,p100-Specific Polyclonal antibody



Catalog Number: 15503-1-AP

Featured Product

5 Publications

Basic Information

Catalog Number:

15503-1-AP

Size:

150ul, Concentration: 300 µg/ml by Nanodrop and 233 µg/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_001077494

GeneID (NCBI):

4791

ENSEMBL Gene ID:

ENSG00000077150

UNIPROT ID:

Q00653

Full Name:

nuclear factor of kappa light polypeptide gene enhancer in B-cells 2 (p49/p100)

Calculated MW:

97 kDa

Observed MW:

52/100 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:20-1:200

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IF, IHC

Species Specificity:

human, mouse

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : A431 cells, HeLa cells, MCF-7 cells, mouse heart tissue

IHC : human testis tissue, human spleen tissue

Background Information

NF-kappa-B is a pleiotropic transcription factor which is present in almost all cell types and is involved in many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFkB1/p105, NFkB1/p50, REL and NFkB2/p52. NFkB2 appears to have dual functions such as cytoplasmic retention of attached NF-kappa-B proteins by p100 and generation of p52 by a cotranslational processing. The proteasome-mediated process ensures the production of both p52 and p100 and preserves their independent function. P52 binds to the kappa-B consensus sequence 5'-GGRNNYYCC-3', located in the enhancer region of genes involved in immune response and acute phase reactions. P52 and p100 are respectively the minor and major form; the processing of p100 being relatively poor. Isoform p49 is a subunit of the NF-kappa-B protein complex, which stimulates the HIV enhancer in synergy with p65. This antibody can bind both p52 and p100 isoforms of NFkB2.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|--------------|-----------|-----------------|-------------|
| Xianhui Yang | 34155950 | Pharm Biol | WB |
| Ha-Yeon Shin | 35159096 | Cancers (Basel) | IHC |
| Jun Wang | 30104686 | Sci Rep | IF, WB |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human testis using 15503-1-AP (NFκB2,p52,p100-Specific antibody) at dilution of 1:50 (under 40x lens).



Immunohistochemical analysis of paraffin-embedded human spleen using 15503-1-AP (NFκB2,p52,p100-Specific antibody) at dilution of 1:50 (under 40x lens).



A431 cells were subjected to SDS PAGE followed by western blot with 15503-1-AP (NFκB2,p52,p100-Specific Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.