For Research Use Only

MKS1 Polyclonal antibody

Catalog Number:16206-1-AP 26 Publications



Basic Information	Catalog Number: 16206-1-AP	GenBank Accession Number: BC010061		Purification Method: Antigen affinity purification	
	Size: GenelD (NCBI):			Recommended Dilutions: WB 1:500-1:2000 IP 0.5-4.0 ug for IP and 1:500-1:2000 for WB	
	150ul, Concentration: 650 µg/ml by Nanodrop and 267 µg/ml by Bradford method using BSA as the standard; Source: Rabbit Isotype:	54903 Full Name: Meckel syndrome, type 1			
					Calculated MW
		559 aa, 65 kDa		IF 1:20-1:200	
		Observed MW:			
		Igu Immunogon Catalog Numbor	65-70 kDa		
	AG9177				
	Applications	Tested Applications:		Positive Controls:	
IF, IHC, IP, WB, ELISA Cited Applications: IF, WB Species Specificity:			WB : mouse b	prain tissue, HEK-293 cells, HeLa cells,	
			mouse uterus tissue, SH-SY5Y cells		
			IP: HEK-293	-293 cells, man liver cancer tissue,	
human, mouse, rat			IHC : human		
Cited Species: human, mouse		IF : nTERT-RPE1 cells and Mouse empryonic fibroblast			
	retrieval may be performed with citrate buffer pH 6.0				
Background Information	MKS1 (Meckel syndrome type 1 protein) is a 559-amino acid protein that contains a conserved B9 domain. It is a component of a large protein complex which localizes to the ciliary transition zone and regulates mammalian ciliogenesis and ciliary membrane composition (PMID: 21725307). MKS1 is required for ciliary structure and function, and is involved in centrosome migration to the apical cell surface during early ciliogenesis (PMID: 17185389; 19515853). Broad tissue expression of the MKS1 gene has been reported (PMID: 16415886). Defects in MKS1 are the cause of Meckel syndrome type 1 (MKS1), an autosomal recessive lethal malformation syndrome characterized by renal cystic dysplasia, central nervous system malformations, and hepatic developmental defects (PMID: 16415886). In addition, defects in MKS1 are also the cause of Bardet-Biedl syndrome type 13 (BBS13) (PMID: 18327255).				
Notable Publications	Author Pu	omed ID Jou	ırnal	Application	
	T Tony Yang 26	365165 Sci	Rep	IF	
	Yunfan Yang 25.	342559 Ce	ll Res	WB	
	Gisela G Slaats 26	490104 J M	led Genet	WB, IF	
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage				
*** 20ul sizes contain 0.1% BSA					

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Mouse Embryonic Fibroblast

Human hTERT-RPE1 Cells

IF result from Dr. Corbit, Kevin. anti-MKS1 (16206-1-AP) marks the transition zone of Human hTERT-RPE1 cells and Mouse embryonic fibroblasts.



mouse brain tissue were subjected to SDS PAGE followed by western blot with 16206-1-AP (MKS1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP Result of anti-MKS1 (IP:16206-1-AP, 3ug; Detection:16206-1-AP 1:1000) with HEK-293 cells lysate 4500ug.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 16206-1-AP (MKS1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 16206-1-AP (MKS1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer using 16206-1-AP (MKS1 antibody) at dilution of 1:50 (under 10x Ìens).



Immunohistochemical analysis of paraffin-embedded human liver cancer using 16206-1-AP (MKS1 antibody) at dilution of 1:50 (under 40x lens).