For Research Use Only

RPS13 Polyclonal antibody

Catalog Number: 16680-1-AP

4 Publications



Basic Information

Catalog Number: 16680-1-AP

GenBank Accession Number:

Purification Method: Antigen affinity purification

Size:

BC006772 GeneID (NCBI):

150ul, Concentration: 600 µg/ml by

6207

Recommended Dilutions: WB 1:1000-1:4000

Nanodrop and 400 µg/ml by Bradford Full Name:

IHC 1:50-1:500 IF 1:50-1:500

method using BSA as the standard;

ribosomal protein S13 Calculated MW:

Rabbit 17 kDa

Isotype: Observed MW: IgG 14 kDa, 17 kDa

Immunogen Catalog Number:

AG10008

Applications

Tested Applications:

IF, IHC, WB, ELISA

Cited Applications:

Positive Controls:

WB: MCF-7 cells, mouse oterus tissue, rat oterus tissue

IHC: human colon cancer tissue,

IF: HeLa cells,

Species Specificity: human, mouse, rat **Cited Species:**

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

RPS13, also named as 40S ribosomal protein S13, belongs to the ribosomal protein S15P family. Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. RPS13 is located in the cytoplasm. It is co-transcribed with two U14 small nucleolar RNA genes, which are located in its third and fifth introns. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

Notable Publications

Author	Pubmed ID	Journal	Application
Keely Oltion	36638793	Cell	WB
Yanrong Zhuang	37369203	Cell	WB
Peng Shuping S	21425412	Stem Cells	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 16680-1-AP (RPS13 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using 16680-1-AP (RP513 antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 16680-1-AP (RPS13 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).