#### For Research Use Only

# GABARAP Polyclonal antibody

Catalog Number: 18723-1-AP

**Featured Product** 

19 Publications

GenBank Accession Number:



**Basic Information** 

Catalog Number:

18723-1-AP BC106748

Size: GeneID (NCBI):

150ul, Concentration: 900 µg/ml by 11337 Nanodrop and 400 µg/ml by Bradford Full Name:

method using BSA as the standard;

GABA(A) receptor-associated protein

Calculated MW: Rabbit 14 kDa Isotype: Observed MW: IgG 14-18 kDa

**Applications** 

**Tested Applications:** 

FC, IF, WB, ELISA **Cited Applications:** 

CoIP, IF, WB

Species Specificity: human, mouse

**Cited Species:** 

human, mouse, monkey

**Positive Controls:** 

WB: HeLa cells, A549 cells, mouse brain tissue, human placenta tissue, mouse pancreas tissue, human heart tissue, mouse skeletal muscle tissue, HuH-7 cells, SH-SY5Y cells, RAW 264.7 cells, mouse liver tissue, rat

**Purification Method:** 

WB 1:1000-1:6000

IF 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

brain tissue

IF: HepG2 cells, Starvation treated HepG2 cells

## **Background Information**

GABARAP (Gamma-aminobutyric acid type A receptor-associated protein), was initially identified as GABAA receptor binding protein and has shown to be implicated in intracellular trafficking of GABAA receptors. Through interacting with a broad rang of proteins, GABARAP has been demonstrated as a multifunctional protein implicated in intracellular vesicle trafficking, apoptosis, and autophagy. The protein level of lipidated form of GABARAP (GABARAP-II) is upregulated during the autophagic process, which makes it as an autophagic marker. This antibody detected endogenous GABARAP protein in lysates from various tissues. It may cross-react with GABARAPL1 due to the high identity of amino acid sequence.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Linyu Sun	34555354	Mol Cell	WB
Svenja Zielke	33111629	Autophagy	WB
Zhen-Hua Chen	27740626	Cell Death Differ	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 18723-1-AP (GABARAP antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human GABARAP (18723-1-AP) and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (-20°C Ethanol) fixed Chloroquine treated HepG2 cells and untreated HepG2 cells using GABARAP antibody (18723-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Rabbit IgG(H+L).