

For Research Use Only

IRF7 Polyclonal antibody

Catalog Number: 22392-1-AP **12 Publications**



Basic Information

Catalog Number: 22392-1-AP	GenBank Accession Number: BC136555	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 500 µg/ml by Nanodrop;	GeneID (NCBI): 3665	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:250-1:1000
Source: Rabbit	UNIPROT ID: Q92985	
Isotype: IgG	Full Name: IRF 7	
Immunogen Catalog Number: AG18059	Calculated MW: 516 aa, 56 kDa	
	Observed MW: 55 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls: WB : HEK-293 cells, IHC : mouse skin tissue,
Cited Applications: WB, IF, IHC	
Species Specificity: human, mouse	
Cited Species: human, Paralichthys olivaceus, mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

IRF-7 (IFN regulatory factor 7) is a member of the IFN regulatory transcription factor (IRF) family. IRF-7 has been shown to play a role in the transcriptional activation of virus-inducible cellular proteins, including IFN beta chain proteins. Inducible expression of IRF-7 is largely restricted to lymphoid tissue. Four transcript variants encoding distinct isoforms (A,B,C and D) have been identified for this (PMID:9786932). The active IRF7A exists as a dimer form ~80 kDa (PMID:11073981). The MW 67-70 kDa has been reported in some papers (PMID:9786932; 22951831). Various posttranslational modifications of IRF7 including phosphorylation, ubiquitination, sumoylation and acetylation are identified (PMID:22951831). This antibody is a rabbit polyclonal antibody raised against the C-terminal 349 amino acid residues of human IRF7 D. The molecular weight of Nonphosphorylated IRF7 cofractionated with the 44-kDa marker, approximating its predicted size. In contrast, phosphorylated IRF7, which migrate more slowly on SDS-PAGE. This phosphorylated IRF7 was consistent with a size of 80 to 90 kDa. (PMID: 11073981)

Notable Publications

Author	Pubmed ID	Journal	Application
Wei Cao	34011520	J Immunol	WB
Zhangchuan Xia	28356387	J Immunol	WB
Xiao-Ming Lyu	29956500	Cancer Med	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with 22392-1-AP (IRF7 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse skin tissue slide using 22392-1-AP (IRF7 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).