For Research Use Only

ATG2B Polyclonal antibody

Catalog Number: 25155-1-AP

Featured Product

14 Publications



Purification Method:

WB 1:500-1:1000

IHC 1:50-1:500

IF 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number:

25155-1-AP BC015016 GeneID (NCBI): Size: 150ul , Concentration: 500 µg/ml by 55102

Nanodrop and 253 μ g/ml by Bradford UNIPROT ID: method using BSA as the standard;

Source: Full Name:

Rabbit ATG2 autophagy related 2 homolog B

Q96BY7

Isotype: (S. cerevisiae) Calculated MW: Immunogen Catalog Number: 2078 aa, 233 kDa AG18412 Observed MW:

240-250 kDa

Applications

Tested Applications: Positive Controls: WB, IF, IHC, ELISA WB: HeLa cells,

Cited Applications: IHC: human pancreas cancer tissue, human stomach

Species Specificity:

human **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Notable Publications

Author	Pubmed ID	Journal	Application
Jay Xiaojun Tan	36071159	Nature	WB
Jee-Eun Lee	34720018	Autophagy	WB
Ying Li	33124742	FASEB J	WB

tissue

IF: HepG2 cells,

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 25155-1-AP (ATG2B antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 25155-1-AP (ATG2B antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



HeLa cells were subjected to SDS PAGE followed by western blot with 25155-1-AP (ATG2B Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using ATG2B antibody (25155-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).