### For Research Use Only

# EOMES/TBR2 Polyclonal antibody

Catalog Number: 28316-1-AP



**Purification Method:** 

WB 1:500-1:1000

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number: GenBank Accession Number:

28316-1-AP NM\_001278182

Size: Genel D (NCBI): 150ul, Concentration: 700 µg/ml by 8320

Nanodrop and 300 µg/ml by Bradford Full Name:

method using BSA as the standard; eomesodermin homolog (Xenopus

Source: laevis)

Rabbit Calculated MW: Isotype: 73 aa

IgG Observed MW:

Immunogen Catalog Number: 70-72 kDa

AG27634

**Applications** 

**Tested Applications:** 

WB,ELISA

Species Specificity: Human, Mouse, rat Positive Controls:

WB: mouse brain tissue,

## **Background Information**

EOMES/TRB2 (also named Eomesodermin/Tbr2) encode T-box transcription factor expressed highly in the intermediate progenitor stage of the developing neuron. During migration of the neurons, radial glia divide and migrate towards the surface of the cerebral cortex, there are three stages of cellular development: radial glia, intermediate progenitors, and postmitotic projection neurons. Radial glia express Pax6, while intermediate progenitor cells express Eomesodermin/Tbr2, and postmitotic projection neurons express Tbr1 (PMID:15634788; PMID:16320258). This process also called neurogenesis, and demonstrated that Eomesodermin/Tbr2 has been implicated in neurodevelopment. According to the mouse model, several articles showed that EOMES/TRB2 KO mice having proliferating cells decreased and having smaller upper cortical layers and a smaller sub ventricular zone in the brain (PMID:18940588). What's more, Eomesodermin/Tbr2 has also been found participating in immune response (PMID:20713880).

#### Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 28316-1-AP (EOMES antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.