For Research Use Only

SMN-Exon7 Monoclonal antibody

Catalog Number:60255-1-lg 1 Publications



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number:

60255-1-lg BC062723 Protein A purification

Size: GeneID (NCBI): CloneNo.: 150ul , Concentration: 1000 µg/ml by 6606 3A8G11

Nanodrop and 543 µg/ml by Bradford method using BSA as the standard; survival of motor neuron 1 telements WB 1:500-1:2000

method using BSA as the standard; survival of motor neuron 1, telomeric WB 1:500-1:2000

Source: Calculated MW: IF 1:50-1:500

 Mouse
 294 aa, 32 kDa

 Isotype:
 Observed MW:

 IgG1
 40 kDa

Immunogen Catalog Number:

AG16615

Applications Tested Applications: Positive Controls:

IF, WB, ELISA WB: HEK-293 cells, HeLa cells, HepG2 cells

Cited Applications:

WB

IF: HepG2 cells,

Species Specificity:

human

Background Information

Spinal muscular atrophy (SMA) is an autosomal recessive neurodegenerative disease characterized by loss of anterior horn cells in the spinal cord and concomitant symmetrical muscle weakness and atrophy (PMID: 16364894). SMA is caused by deletion or mutations of the survival motor neuron (SMN1) gene. SMA patients lack a functional SMN1 gene, but they possess an intact SMN2 gene, which though nearly identical to SMN1, is only partially functional (PMID: 17355180). A large majority of SMN2 transcripts lack exon 7, resulting in production of a truncated, less stable SMN protein (PMID: 10369862). The level of SMN protein correlates with phenotypic severity of SMA. This antibody, 60255-1-lg, raised against the C-terminal region (275-294aa) encoded by the exon 7.

Notable Publications

Author	Pubmed ID	Journal	Application
Mandana Arbab	36996170	Science	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with 60255-1-1g (SMN-Exon7 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells using 60255-1-Ig (SMN-Exon7 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).