

For Research Use Only

Anti-Human CD28 (CD28.2)

Catalog Number: 65099-1-Ig **3 Publications**



Basic Information

Catalog Number: 65099-1-Ig	GenBank Accession Number: BC093698	Purification Method: Affinity purification
Size: 100ug, 0.5 mg/ml	GeneID (NCBI): 940	CloneNo.: CD28.2
Source: Mouse	Full Name: CD28 molecule	
Isotype: IgG1, kappa	Calculated MW: 220 aa, 25 kDa	

Applications

Tested Applications:
FC

Cited Applications:
Cell treatment, IHC

Species Specificity:
Human

Cited Species:
human

Background Information

CD28 (T-cell-specific surface glycoprotein CD28), also known as T44 and Tp44, is a 44 kD disulfide-linked homodimeric type I glycoprotein (PMID: 2162180). It is a member of the immunoglobulin superfamily and is expressed on most T lineage cells, NK cell subsets, and plasma cells (PMID: 2162180, 8386518). CD28 may affect in vivo immune responses by functioning both as a cell adhesion molecule linking B and T lymphocytes and as the surface component of a novel signal transduction pathway (PMID: 2162180, 3021470). CD28 binds both CD80 and CD86 with a highly conserved motif MYPPY in the CDR3-like loop (PMID: 15696168, 7964482). CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death (PMID: 1348520).

Notable Publications

Author	Pubmed ID	Journal	Application
Wenbo Zhao	36309750	J Transl Med	
Ya-Nan Pi	34852825	Cancer Cell Int	IHC
Guojun Yao	37354488	Aging (Albany NY)	Cell treatment

Storage


Storage:
Store at 2-8°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.09% sodium azide.


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ human PBMCs were surface co-stained with CL647 Plus Anti-Human CD3 and 0.2 ug Anti-Human CD28 (65099-1-Ig, Clone:CD28.2) with CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000. Cells were not fixed. Lymphocytes were gated.



1X10⁶ human PBMCs were surface stained with CL647 Plus Anti-Human CD3 and 0.2 ug Anti-Human CD28 (65099-1-Ig, Clone:CD28.2) with CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 or unstained. Cells were not fixed. Lymphocytes were gated.