For Research Use Only

Alpha Galactosidase A Monoclonal antibody



Catalog Number:66121-1-Ig

Featured Product

2 Publications

49 kDa

Basic Information

Catalog Number: GenBank Accession Number: 66121-1-lg BC002689

ze: GeneID (NCBI):

150ul , Concentration: 3900 µg/ml by 2717 Nanodrop and 2387 µg/ml by Bradford $_{\hbox{Full Name}}$

method using BSA as the standard; galactosidase, alpha

Source: Calculated MW:
Mouse 49 kDa
Isotype: Observed MW:

lgG2a

Immunogen Catalog Number:

AG7505

Purification Method: Protein A purification

CloneNo.: 2B2C5

Recommended Dilutions:

WB 1:3000-1:10000 IHC 1:20-1:200 IF 1:20-1:200

Applications

Tested Applications:

IF, IHC, WB, ELISA

Cited Applications:

WB

Species Specificity:

human
Cited Species:
human

human Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen

retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, HEK-293 cells, HepG2 cells, human

kidney tissue, LNCaP cells, A431 cells

IHC: human kidney tissue, human liver tissue

IF: HepG2 cells,

Background Information

GLA, also named as Melibiase and Alpha-galactosidase A, belongs to the glycosyl hydrolase 27 family. It hydrolyzes terminal, non-reducing alpha-D-galactose residues in alpha-D-galactosides, including galactose oligosaccharides, galactomannans and galactolipids. Fabry disease is an X-linked lysosomal storage disorder resulting from the deficient activity of GLA. Enzyme replacement therapy (ERT) with GLA is currently the most effective therapeutic strategy for patients with Fabry disease, a lysosomal storage disease.

Notable Publications

Author	Pubmed ID	Journal	Application
Sebastian Boland	36207292	Nat Commun	WB
Chenxu Guo	38328234	bioRxiv	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




WB result of GLA antibody (66121-1-Ig, 1:2000) with si-Control and si-GLA transfected HEK293 cells..




Various lysates were subjected to SDS PAGE followed by western blot with 66121-1-1g (Alpha galactosidase A antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 66121-1-Ig (Alpha galactosidase A antibody at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 66121-1-Ig (Alpha galactosidase A antibody at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of HepG2 cells using 66121-1-1g (Alpha galactosidase A antibody) at dilution of 1:50 and Rhodamine-Goat anti-Mouse IgG.