

For Research Use Only

CD80/B7-1 Monoclonal antibody

Catalog Number: 66406-1-Ig **32 Publications**



Basic Information

Catalog Number: 66406-1-Ig	GenBank Accession Number: BC042665	Purification Method: Protein A purification
Size: 150ul , Concentration: 1500 µg/ml by Nanodrop;	GeneID (NCBI): 941	CloneNo.: 1E2F10
Source: Mouse	UNIPROT ID: P33681	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500
Isotype: IgG1	Full Name: CD80 molecule	
Immunogen Catalog Number: AG5615	Calculated MW: 33 kDa	
	Observed MW: 68 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls: WB : Raji cells, DC2.4 cells, Daudi cells IHC : human tonsillitis tissue,
Cited Applications: WB, IF, FC, IHC	
Species Specificity: human, mouse, pig	
Cited Species: human, rat, mouse, rabbit, pig	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

CD80 (also known as B7-1) is a type I membrane protein that is a member of the immunoglobulin superfamily, with an extracellular immunoglobulin constant-like domain and a variable-like domain required for receptor binding. It is expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, monocytes, and macrophages. CD80 is the receptor for the proteins CD28 and CTLA-4 found on the surface of T-cells. It is involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is induced by the binding of CD28, binding to CTLA-4 has opposite effects and inhibits T-cell activation. CD80 also acts as a cellular attachment receptor for adenovirus subgroup B. (PMID: 7545666; 12015893; 16920215)

Notable Publications

Author	Pubmed ID	Journal	Application
Yan Yang	34582523	Biomater Sci	FC
Chenxia Juan	30243953	Int J Biochem Cell Biol	WB
Yuanhang Li	30411299	Pathol Oncol Res	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Raji cells were subjected to SDS PAGE followed by western blot with 66406-1-Ig (CD80 / B7-1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66406-1-Ig (CD80 / B7-1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66406-1-Ig (CD80 / B7-1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).