

For Research Use Only



SARS-CoV-2 Nucleocapsid Phosphoprotein Recombinant antibody

Catalog Number: 80027-1-RR

Basic Information

Catalog Number: 80027-1-RR	GenBank Accession Number: NC_045512	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 43740575	CloneNo.: 8C20
Source: Rabbit	Full Name: COVID-19 N Protein	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG		
Immunogen Catalog Number: AG30676		

Applications

Tested Applications: WB, ELISA	Positive Controls:
Species Specificity: virus	WB : Eukaryotic nucleocapsid phosphoprotein, Ag30676

Background Information

The nucleocapsid (N) protein has multiple functions including formation of nucleocapsids, signal transduction virus budding, RNA replication, and mRNA transcription. N protein is an important antigen for coronavirus, and it is normally highly conserved, with a molecular weight of about 50 kDa. It can be used as a marker in diagnostic assays due to its high immunogenicity (PMID: 32416961, PMID: 32235387). A sandwich ELISA for COVID-19 N Protein can be assembled by using 80027-1-RR as capture antibody and conjugated 80026-1-RR for detection.

Storage


Storage:
Store at -20°C.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




E.coli expressed SARS-CoV-2 Nucleocapsid Phosphoprotein (Cat.NO. Ag30676) was subjected to SDS-PAGE followed by western blot with 80027-1-RR and 80026-1-RR at various work concentration.



Eukaryotic expressed SARS-CoV-2 Nucleocapsid Phosphoprotein was subjected to SDS-PAGE followed by western blot with 80027-1-RR and 80026-1-RR at various work concentration.



Indirect ELISA was carried out by coating eukaryotic expressed N protein at 70 ng/well followed by blocking and adding serial diluted primary antibody 80026-1-RR and 80027-1-RR respectively. Signal was developed with TMB and stopped by H₂SO₄. Signal strength was measured by absorbance at 450 nm.



Sandwich ELISA was carried out by coating 80027-1-RR at 80 ng/well followed by blocking and adding different concentration of eukaryotic expressed N protein (0.5-1000 ng/mL). HRP-conjugated 80026-1-RR was used at 1 µg/mL for detection. Signal was developed with TMB and stopped by H₂SO₄. Signal strength was measured by absorbance at 450 nm.