

Catalog Number: HZ-1011

## **Data Sheet**



## HumanKine® TGF beta 1 (Recombinant Human)

**Animal Component-Free** 


**Human cell expressed** 

Tag-Free


**Endotoxin Free** 

Product Description				
Animal-free Recombinant Human TGF beta 1 is expressed from human 293 cells as a mature disulfide-linked homodimer with an apparent molecular mass of 25 kDa. Tests indicate that TGF beta 1 is as efficient at inducing IL-17 expression in Th17 cells, as cytokine purified from human platelets. This product is produced in a human cell expression system with serum-free, chemically defined media.				
Alternative Names	lternative Names CED, DPD1, LAP, TGF B1, TGF beta 1, tgf beta1, TGFB, TGFB1, TGFbeta, tgf-beta1			
Source	Source Human Embryonic Kidney cells (HEK293). HEK293-derived TGF beta 1 protein			
Species Reactivity human,mouse,goat,pig,rat				

Specifications					
Test	Method	Specification			
Activity	Dose-dependent inhibition of IL-4 induced proliferation of mouse HT-2 cells (BALB/c spleen activated by sheep erythrocytes in the presence of IL-2)	Typically ≤ 0.5 ng/mL EC50			
Molecular Mass	SDS-PAGE	13 kDa reduced, 25 kDa non-reduced, homodimer, non-glycosylated			
Purity	SDS-PAGE	>95%			
Endotoxin	LAL	<1 EU/µg			



Recombinant human TGF beta 1 (HZ-1011) inhibits IL-4 induced proliferation of the HT-2 mouse cell line. HT-2 cells are Balb/c spleen cells activated by sheep erythrocytes in the presence of IL-2. Cell number was quantitatively assessed by PrestoBlue® cell viability reagent. HT-2 cells were treated with increasing concentrations of recombinant TGF beta 1 for 72 hours. The EC50 was



www.ptglab.com

Document #: FR-QA118-101 Rev 0
Data Sheet Version #: 1

Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com

Preparation				
Shipping Temperature	ambient temperature			
Formulation	50 mM NaOAc pH 3.7, See Certificate of Analysis for details			
Reconstitution  Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile 4 mM HCl containing endotoxin-free recombinant human serum albumin (HSA).				

Stability and Storage	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
	Lyophilized	Room Temperature	2 weeks
	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
	Avoid repeated freeze-thaw cycles.		

www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com