

HumanKine® IL-10 (Recombinant Human)



Animal Component-Free	Human cell expressed	Tag-Free	Endotoxin Free
-----------------------	----------------------	----------	----------------

Product Description


Animal-free Recombinant Human IL-10 is expressed in human 293 cells as a non-covalent homodimeric glycoprotein with an apparent molecular mass of 17 kDa. This cytokine is produced in a serum-free, chemically defined media. IL-10 is produced by a variety of mammalian cell types which includes T cells, B cells, macrophages, and keratinocytes monocytes. It is an anti-inflammatory cytokine with multiple, pleiotropic effects in inflammation and immunoregulation. Glycosylation contributes to stability in cell growth media and other applications. The cytokine is greater than 95% pure.

Alternative Names	CSIF, IL 10, IL10, IL10A, interleukin 10, TGIF
Source	Human Embryonic Kidney cells (HEK293). HEK293-derived IL-10 protein
Species Reactivity	human,mouse,pig

Specifications


Test	Method	Specification
Activity	Dose-dependent proliferation of the MC/9 (mouse mast cell) cell line	0.18-2.0 ng/mL EC50
Molecular Mass	SDS-PAGE	19 kDa reduced, 17 kDa non-reduced, non-disulfide bonded homodimer, glycosylated
Purity	SDS-PAGE	>95%
Endotoxin	LAL	<1 EU/μg

Activity Data



Recombinant human IL-10 (HZ-1145) induces dose-dependent proliferation of the MC/9 (mouse mast cell) cell line. Cell number was quantitatively assessed by PrestoBlue® cell viability reagent. MC/9 cells were treated with increasing concentration of recombinant IL-10 for 48 hours. The EC50 was determined using a 4-parameter non-linear regression model. The EC50

SDS-PAGE



Preparation	
Shipping Temperature	ambient temperature
Formulation	10mM Tris-HCl pH 7.4 + 150mM NaCl, See Certificate of Analysis for details
Reconstitution	Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile 1xPBS pH 7.4 containing 0.1% endotoxin-free recombinant human serum albumin (HSA).

Stability and Storage	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
	Lyophilized	Room Temperature	2 weeks
	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
Avoid repeated freeze-thaw cycles.			

www.ptglab.com

Document #: FR-QA118-101 Rev 0
Data Sheet Version #: 1

Proteintech Group, Inc.
5500 Pearl Street, Suite 400 Rosemont, IL 60612
t: 1-888-478-4522; f: 1-312-455-8408
Email: proteintech@ptglab.com