

Species Reactivity

Test

Catalog Number: HZ-1218

Data Sheet



Animal Component-Free Human cell expressed Tag-Free

Endotoxin Free

Product Description Animal-free Recombinant Human FGF-4 is expressed in human 293 cells as a monomeric glycoprotein with an apparent molecular mass of 17 and 27 kDa. This cytokine is produced in a human cell expression system with serum-free, chemically defined media. Authentic glycosylation contributes to stability in cell growth media and other applications. FGF-4 plays an important physiologically role in the self-renewal of human embryonic stem cells. It also promotes stem cell proliferation. Furthermore, FGF-4 may also aid differentiation which additionally relies on context and concentration. FGF 4, FGF4, FGF-4, fibroblast growth factor 4, HBGF 4, HST, HST 1, HSTF 1, HSTF1, K FGF, KFGF, KS3, Transforming protein KS3 **Alternative Names** Human Embryonic Kidney cells (HEK293). HEK293-derived FGF-4 protein Source

human,mouse

Specifications Method Specification Dose-dependent stimulation of the proliferation of the Balb/c 3T3 cell line. Dose-≤ 1.25 ng/mL EC50 in Balb/c 3T3 cells. 6-30 ng/mL

Activity Dependent Proliferation of the NIH3T3 cell line in defined media EC50 in NIH 3T3 cells in defined media 17 and 27 kDa reduced and non-reduced, monomer, Molecula SDS-PAGE glycosylated Mass SDS-PAGE >95% Purity <1 EU/µg Endotoxir LAL



www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1

Proteintech Group, Inc. 5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com

Preparation			
Shipping Temperature	ambient temperature		
Formulation	1x PBS, See Certificate of Analysis for details		
Reconstitution Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile 1x PBS pH 7.4.			

Stability and Storage	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
	Lyophilized	Room Temperature	2 weeks
	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
	Avoid repeated freeze-thaw cycles.		

www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc. 5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com