



IHCeasy Human IgG kappa chain Ready-To-Use IHC Kit

Catalog Number: KHC0072

General Information

Sample type: FFPE tissue Cited sample type: Reactivity: Human Cited Reactivity: Assay type: Immunohistochemistry Primary antibody type: Mouse Monoclonal

Secondary antibody type: Polymer-HRP-Goat anti-Mouse

Kit Component

Component	Size	Concentration
Antigen Retrieval Buffer	100 mL	50×
Washing Buffer	100 mL ×2	20×
Blocking Buffer	5 mL	RTU
Primary Antibody	5 mL	RTU
Secondary Antibody	5 mL	RTU
Chromogen Component A	0.2 mL	RTU
Chromogen Component B	4 mL	RTU
Signal Enhancer	5 mL	RTU
Counter Staining Reagent	5 mL	RTU
Mounting Media	5 mL	RTU
Control Slide	1 slide (Optional)	FFPE
Datasheet	1 Copy	
Manual	1 Copy	

Storage Instructions

All the reagents are stored at 2-8°C. The kit is stable for 6 months from the date of receipt.


Background

All immunoglobulins consists of 2 heavy chains and 2 light chains. The light chains are designated kappa and lambda. In human and most mammals, Ig-molecules containing kappa light chains are slightly more frequent than molecules containing lambda. Abnormal kappa/lambda ratio implies neoplasms of B-lymphocytes (lymphomas and leukemias). It has also been reported that free light chains (FLC) kappa/lambda may be specific and sensitive for the diagnosis of type 2 diabetes mellitus (T2D).

Synonyms

Human IgG Kappa chain, IgK

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using KHC0072 (Human IgG Kappa chain IHC Kit).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using KHC0072 (Human IgG Kappa chain IHC Kir)