

## GFP-Booster for Immunofluorescence Detection of GFP-Fusion Proteins

For the immunofluorescence detection of GFP-fusion proteins in fixed cells.

---

- 1. Product** The GFP-Booster Alexa Fluor® 568 is an anti-GFP Nanobody coupled to Alexa Fluor® 568.
- 2. Introduction** Green fluorescent protein (GFP) and its variants are widely used to study protein localization and dynamics in cells. However, photo-stability and quantum efficiency of GFP are often not sufficient for e.g. super-resolution microscopy (such as 3D-SIM or dSTORM) and for fixed cell samples. In addition, many cell biological methods such as BrdU-staining, EdU-Click-iT™ treatment or fluorescent *in situ* hybridization result in disruption of the GFP signal. The GFP-Booster reactivates, enhances, and stabilizes the GFP-signal.
- 3. Properties**
- |                       |                                                                                                                                               |
|-----------------------|-----------------------------------------------------------------------------------------------------------------------------------------------|
| Product size          | gb2AF568-10: 10 µL<br>gb2AF568-50: 50 µL                                                                                                      |
| Format                | Alpaca single domain antibody, Nanobody or V <sub>H</sub> H; monovalent                                                                       |
| Target/ Specificity   | GFP and GFP variants. See <a href="http://www.chromotek.com">www.chromotek.com</a> for a list of recognized GFP variants.                     |
| Conjugate             | Site-directed conjugation to Alexa Fluor 568                                                                                                  |
| Excitation/ Emission  | Excitation max: 578 nm, Emission max: 603 nm                                                                                                  |
| DOL                   | 2 fluorophores per Nanobody                                                                                                                   |
| Purity                | Recombinantly expressed and purified                                                                                                          |
| Form                  | Buffered aqueous solution                                                                                                                     |
| Storage buffer        | 10 mM HEPES pH 7.0, 500 mM NaCl, 5 mM EDTA,<br>Preservative: 0.09% Sodium azide, Safety datasheet (SDS): <b>Sodium azide SDS</b>              |
| Concentration         | 0.5 g/L                                                                                                                                       |
| Stability and storage | Shipped at ambient temperature. Store at -20°C/-4°F. Avoid freeze-thaw cycles. Aliquot upon arrival. Protect from light. Stable for 6 months. |
- 4. Protocol**
- Fixation:** Fix cells seeded on coverslips in 3.7% formaldehyde in PBS for 10 min at room temperature.  
*Note: Always prepare a fresh formaldehyde dilution.*  
*Note: Alternatively, use methanol for fixation: Apply ice-cold 100% methanol to cells for 3 min, wash as in step 2 and proceed directly with step 5 of this protocol.*
  - Wash samples three times with PBS (Phosphate Buffered Saline). Do not store fixed cells.
  - Permeabilization:** Add PBS containing 0.5% Triton X-100 to samples and incubate for 5 min at room temperature.
  - Wash samples twice with PBS.
  - Blocking:** Add 4% BSA in PBS to samples and incubate for 10 min at room temperature.
  - GFP-Booster incubation:** Dilute GFP-Booster 1:500 – 1:1,000 in blocking buffer and incubate for 1 h at room temperature. Optimal dilution is application-dependent and should be determined.  
*Note: For multiplexing protocols, you can combine GFP-Booster with any other antibody.*
  - Wash samples three times for 5-10 min in PBS.
  - If required, counter stain with DNA fluorescent dyes, e.g. DAPI in PBS.
  - Mounting:** Rinse sample shortly in water to prevent formation of salt crystals. Mount

in VectaShield (Vector Labs) or other mounting media with anti-fading agents and seal mounted coverslips with clear nail polish.

**Suggested buffer composition**

Buffer	Composition
Blocking buffer	4% BSA (w/v) in PBS
Fixation buffer	3.7% formaldehyde in PBS
Permeabilization buffer	PBS; 0.5% Triton X-100
Wash buffer	PBS

**5. Support/  
Troubleshooting**

Please refer to our FAQ section at [www.chromotek.com](http://www.chromotek.com) or contact [support@chromotek.com](mailto:support@chromotek.com)

*Only for research applications, not for diagnostic or therapeutic use.*

ChromoTek is a registered trademark of ChromoTek GmbH. Nanobody is a registered trademark of Ablynx, a Sanofi company. Alexa Fluor is a registered trademark of Life Technologies Corporation, a part of Thermo Fisher Scientific Inc.