For Research Use Only

Marcks Polyclonal antibody

Catalog Number:10004-2-lg Featured Product 14 Publications



Basic Information	Catalog Number: 10004-2-lg	GenBank Accession Number: BC046601	Purification Method: Protein A purification			
	Size:	GenelD (NCBI):	Recommended Dilutions:			
	150ul , Concentration: 2000 ug/ml by		WB 1:500-1:1000			
	Nanodrop and 947 ug/ml by Bradford method using BSA as the standard; Source: Rabbit Isotype: IgG	UNIPROT ID:	IHC 1:50-1:500			
		P26645				
		Full Name: myristoylated alanine rich protein kinase C substrate				
				Calculated MW: 39 kDa		
				Observed MW:		
		88 kDa				
		Applications	Tested Applications:	Positive Controls:		
			WB, IHC, ELISA	WB : mouse	WB: mouse brain tissue,	
Cited Applications: WB, IHC, IF, IP	IHC : mouse pancreas tissue,					
Species Specificity: mouse						
Cited Species: mouse, rat, bovine						
Note-IHC: suggested antigen ro TE buffer pH 9.0; (*) Alternativ retrieval may be performed wa buffer pH 6.0	vely, antigen					
Background Information	apparent molecular weights were fou	ue distribution. The calculated mole Ind at 68 kDa to 90 kDa in different nd membrane recycling. The anti-M	ecular weight of MARCKS is 39 kDa and i species. This protein is a known to IARCKS antibody is raised against purifi			
	activation. It shows a ubiquitous tissu apparent molecular weights were fou regulated cytoskeleton remodeling a MARCKS protein from mouse brain us	ue distribution. The calculated mole Ind at 68 kDa to 90 kDa in different nd membrane recycling. The anti-N ing a method similar to that previo	ecular weight of MARCKS is 39 kDa and i species. This protein is a known to IARCKS antibody is raised against purifi- usly reported for MacMARCKS			
	activation. It shows a ubiquitous tissu apparent molecular weights were fou regulated cytoskeleton remodeling a MARCKS protein from mouse brain us Author Pub	ue distribution. The calculated mole and at 68 kDa to 90 kDa in different nd membrane recycling. The anti-M ing a method similar to that previo med ID Journal	ecular weight of MARCKS is 39 kDa and i species. This protein is a known to IARCKS antibody is raised against purifi- usly reported for MacMARCKS Application			
	activation. It shows a ubiquitous tissu apparent molecular weights were fou regulated cytoskeleton remodeling a MARCKS protein from mouse brain us Author Pub Zhenzhen Chen 295	ue distribution. The calculated mole and at 68 kDa to 90 kDa in different nd membrane recycling. The anti-N ing a method similar to that previo med ID Journal 188366 J Biol Chem	ecular weight of MARCKS is 39 kDa and i species. This protein is a known to IARCKS antibody is raised against purifi- usly reported for MacMARCKS Application WB,IHC			
	activation. It shows a ubiquitous tissu apparent molecular weights were fou regulated cytoskeleton remodeling a MARCKS protein from mouse brain us Author Pub Zhenzhen Chen 295 Qing-Yan Zhang 358	ue distribution. The calculated mole and at 68 kDa to 90 kDa in different nd membrane recycling. The anti-M ing a method similar to that previo med ID Journal 188366 J Biol Chem	ecular weight of MARCKS is 39 kDa and i species. This protein is a known to IARCKS antibody is raised against purifi- usly reported for MacMARCKS Application WB,IHC			
Background Information Notable Publications	activation. It shows a ubiquitous tissu apparent molecular weights were fou regulated cytoskeleton remodeling a MARCKS protein from mouse brain us Author Pub Zhenzhen Chen 295 Qing-Yan Zhang 358	e distribution. The calculated mole and at 68 kDa to 90 kDa in different nd membrane recycling. The anti-N ing a method similar to that previo med ID Journal 188366 J Biol Chem 150487 Mol Cell Endocrin	ecular weight of MARCKS is 39 kDa and i species. This protein is a known to IARCKS antibody is raised against purifi usly reported for MacMARCKS Application WB,IHC ol WB			
	activation. It shows a ubiquitous tissu apparent molecular weights were fou regulated cytoskeleton remodeling a MARCKS protein from mouse brain us Author Pub Zhenzhen Chen 295 Qing-Yan Zhang 358 Mingyue Zhong 357 Storage: Storage: Storage Buffer:	e distribution. The calculated mole and at 68 kDa to 90 kDa in different nd membrane recycling. The anti-M ing a method similar to that previo med ID Journal 188366 J Biol Chem 150487 Mol Cell Endocrin 126797 Food Funct er shipment.	ecular weight of MARCKS is 39 kDa and i species. This protein is a known to IARCKS antibody is raised against purifi- usly reported for MacMARCKS Application WB,IHC ol WB			
Notable Publications	activation. It shows a ubiquitous tissu apparent molecular weights were fou regulated cytoskeleton remodeling a MARCKS protein from mouse brain us Author Pub Zhenzhen Chen 295 Qing-Yan Zhang 358 Mingyue Zhong 357 Storage: Storage:	e distribution. The calculated mole and at 68 kDa to 90 kDa in different nd membrane recycling. The anti-M ing a method similar to that previo med ID Journal 38366 J Biol Chem 350487 Mol Cell Endocrin 26797 Food Funct er shipment. % glycerol pH 7.3.	ecular weight of MARCKS is 39 kDa and i species. This protein is a known to IARCKS antibody is raised against purifi usly reported for MacMARCKS Application WB,IHC ol WB			


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffinembedded mouse pancreas tissue slide using 10004-2-1g (Marcks antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse pancreas tissue slide using 10004-2-1g (Marcks antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse brain tissue were subjected to SDS PAGE followed by western blot with 10004-2-Ig (Marcks antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.