

PLK1

Polyclonal ANTIBODY

Catalog Number: 10305-1-AP

1 Publications

Basic Information

Catalog Number:
10305-1-AP

Size:
24 µg/150 µl

Source:
Rabbit

Isotype:
IgG

Purification Method:
Antigen affinity purification

Immunogen Catalog Number:
AG0284

GenBank Accession Number:
BC002369

GeneID (NCBI):
5347

Full Name:
polo-like kinase 1 (Drosophila)

Calculated MW:
68 kDa

Observed MW:
62 kDa

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:50-1:500

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

WB

Species Specificity:

human,mouse,rat

Positive Controls:

WB : U2OS cells; MDA-MB-453s cells, COLO 320 cells, A375 cells, mouse testis tissue

IHC : human testis tissue; human cervical cancer tissue, human placenta tissue

Background Information

PLK1(Polo-like kinase 1), also named as PLK and STPK13, belongs to the protein kinase superfamily, Ser/Thr protein kinase family and CDC5/Polo subfamily. PLK1 is a Serine/threonine-protein kinase that performs several important functions throughout M phase of the cell cycle, including the regulation of centrosome maturation and spindle assembly, the removal of cohesins from chromosome arms, the inactivation of APC/C inhibitors, and the regulation of mitotic exit and cytokinesis. The localization of Polo changes during the cell cycle. In interphase, Polo shows a mostly diffuse cytoplasmic localization. Polo becomes concentrated on centrosomes from early prophase and appears on centromeres or kinetochores from late prophase, before nuclear envelope breakdown (NEB). After anaphase onset, Polo relocates to the central spindle and remains enriched at the midbody ring in late stages of cytokinesis (PMID: 29167465).

Notable Publications

Author	Pubmed ID	Journal	Application
Yixi Xu	29106372	Elife	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



U2OS cells were subjected to SDS PAGE followed by western blot with 10305-1-AP (PLK1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours



Immunohistochemistry of paraffin-embedded human testis tissue slide using 10305-1-AP (PLK1 antibody) at dilution of 1:200 (under 10x lens) heat mediated antigen retrieved with Tris-EDTA buffer(pH9).



Immunohistochemistry of paraffin-embedded human testis tissue slide using 10305-1-AP (PLK1 antibody) at dilution of 1:200 (under 40x lens) heat mediated antigen retrieved with Tris-EDTA buffer(pH9).