

For Research Use Only

DKK3 Polyclonal antibody

Catalog Number:10365-1-AP

Featured Product

24 Publications



Basic Information

Catalog Number: 10365-1-AP	GenBank Accession Number: BC007660	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 650 ug/ml by Nanodrop and 393 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 27122	Recommended Dilutions: WB 1:500-1:2000 IHC 1:250-1:1000
Source: Rabbit	UNIPROT ID: Q9UBP4	
Isotype: IgG	Full Name: dickkopf homolog 3 (Xenopus laevis)	
Immunogen Catalog Number: AG0417	Calculated MW: 38 kDa	
	Observed MW: 38-55 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls: WB : HUVEC cells, MCF-7 cells IHC : human liver cancer tissue,
Cited Applications: WB, IHC, IF	
Species Specificity: human, mouse	
Cited Species: human, mouse, rat, canine	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

DKK3, also named as REIC, belongs to the dickkopf family. It antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt and by forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6. DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as antero-posterior axial patterning, limb development, somitogenesis and eye formation. In the adult, Dkks are implicated in bone formation and bone disease, cancer and Alzheimer's disease. The adenovirus carrying REIC/Dkk-3 (Ad-REIC) exhibits a potent tumor-specific cell-killing function for various human cancers. It has also become evident that some human cancers are resistant to Ad-REIC-induced apoptosis. The DKK3 is a Glycosylation protein with MW about 55 kDa or 38-43 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Lulin Huang	36115892	Sci China Life Sci	IF
Laury Amelia Roa Fuentes	36303276	Eur J Oral Sci	IHC
Bo Liu	31772644	Exp Ther Med	WB,IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



WB result of DKK3 antibody (10365-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-DKK3 transfected MCF-7 cells.



HUVEC cells were subjected to SDS PAGE followed by western blot with 10365-1-AP (DKK3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 10365-1-AP (DKK3 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).