

For Research Use Only

FMNL1 Polyclonal antibody

Catalog Number: 10395-1-AP



Basic Information

Catalog Number: 10395-1-AP	GenBank Accession Number: BC001710	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 245 µg/ml by Nanodrop;	GeneID (NCBI): 752	Recommended Dilutions: WB 1:200-1:500 IHC 1:50-1:500 IF 1:200-1:800
Source: Rabbit	Full Name: formin-like 1	
Isotype: IgG	Calculated MW: 160 kDa	
Immunogen Catalog Number: AG0594	Observed MW: 150-160 kDa	

Applications

Tested Applications: IF, IHC, WB, ELISA	Positive Controls: WB : Raji cells, Jurkat cells, HeLa cells IHC : human small intestine tissue, IF : U2OS cells, Saos-2 cells
Species Specificity: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

FMNL1, also named as C17orf1, C17orf1B and FMNL, belongs to the formin homology family. It may play a role in the control of cell motility and survival of macrophages. FMNL protein may be involved in anti-apoptosis mechanisms. It is overexpressed in lymphoid malignances and is found associated with Akt. The MW of FMNL1 is about 150-160 kDa. (PMID: 16494944)

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Raji cells were subjected to SDS PAGE followed by western blot with 10395-1-AP (FMNL1 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human small intestine tissue slide using 10395-1-AP (FMNL1 antibody) at dilution of 1:200 (under 10x lens).



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using FMNL1 antibody (10395-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Methanol) fixed Saos-2 cells using FMNL1 antibody (10395-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Jurkat cells were subjected to SDS PAGE followed by western blot with 10395-1-AP (FMNL1 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.