For Research Use Only

GATA3 Polyclonal antibody

Catalog Number: 10417-1-AP

Featured Product

29 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Antigen affinity purification

10417-1-AP

BC003070 GeneID (NCBI):

Recommended Dilutions:

150ul , Concentration: 500 $\mu g/ml$ by

WB 1:500-1:1000

IHC 1:50-1:500

Nanodrop and 313 µg/ml by Bradford Full Name:

2625

method using BSA as the standard;

GATA binding protein 3

Rabbit

Calculated MW: 443 aa, 48 kDa

Isotype: IgG

Observed MW:

45-50 kDa

Immunogen Catalog Number:

AG0174

Tested Applications:

IHC, WB, ELISA

Cited Applications:

ChIP, WB

Species Specificity:

human

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Applications

Positive Controls:

WB: MCF-7 cells, HeLa cells

IHC: human breast cancer tissue.

Background Information

GATA3 is a transcriptional factor that binds to the enhancer elements of all 4T-cell antigen receptor genes. GATA3 is highly expressed in naive, freshly activated cells and Th2 lineage cells and thought to be necessary and sufficient for Th2 cytokine gene expression. GATA3 and TBET interfer with each other to regulate gene expression. This is a rabbit polyclonal antibody raised against part of C-terminus of human GATA3.

Notable Publications

Author	Pubmed ID	Journal	Application
Shi-Jie Bi	34508802	J Ethnopharmacol	
Mengheng Wang	34539800	Evid Based Complement Alternat Med	
Weiwei Sheng	29072694	Cell Death Dis	WB

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10417-1-AP (GATA3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



MCF-7 cells were subjected to SDS PAGE followed by western blot with 10417-1-AP (GATA3 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.