

For Research Use Only

P53 Polyclonal antibody

Catalog Number: 10442-1-AP

Featured Product

1110 Publications



Basic Information

Catalog Number:

10442-1-AP

Size:

150ul, Concentration: 600 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0698

GenBank Accession Number:

BC003596

GeneID (NCBI):

7157

UNIPROT ID:

P04637

Full Name:

tumor protein p53

Calculated MW:

44 kDa

Observed MW:

53 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:5000-1:50000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC: 1:250-1:1000

IF/ICC: 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP, ChIP, RIP

Species Specificity:

human, rat

Cited Species:

human, rat, pig, rabbit, monkey, chicken, zebrafish, sheep, goat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A431 cells, SMMC-7721 cells, MCF-7 cells, rat colon tissue, HEK-293 cells, HT-29 cells, HepG2 cells

IP: HEK-293 cells,

IHC: human ovary cancer tissue,

IF/ICC: A431 cells,

Background Information

TP53, also known as P53 and NY-CO-13, belongs to the p53 family and has 9 isoforms. In SDS-Page, the observed molecular weight is about 53 kDa. TP53 acts as a tumor suppressor in many tumor types, including growth arrest or apoptosis depending on the physiological circumstances and cell types. It is involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. TP53 Localizes in the nucleus in most cells but found in the cytoplasm in some cells. (PMID: 26166714; PMID: 25225161)

Notable Publications

Author	Pubmed ID	Journal	Application
Sushan Yang	36180831	Cell Mol Biol Lett	WB
Xiaoyan Liu	31574948	Int J Mol Sci	WB
Shanshan Hu	33134183	Front Oncol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10442-1-AP (P53 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




WB result of P53 antibody (10442-1-AP; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-P53 transfected A431 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 10442-1-AP (P53 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




IP result of anti-P53 (IP:10442-1-AP, 4ug; Detection:10442-1-AP 1:20000) with HEK-293 cells lysate 1560 ug.



Immunohistochemical analysis of paraffin-embedded human ovary cancer tissue slide using 10442-1-AP (P53 antibody) at dilution of 1:500 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using P53 antibody (10442-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using P53 antibody (10442-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).