## For Research Use Only

## STMN2 Polyclonal antibody

Catalog Number: 10586-1-AP

Featured Product

29 Publications

GenBank Accession Number:



**Purification Method:** 

WB 1:1000-1:6000

protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

**Basic Information** 

**Applications** 

Catalog Number:

10586-1-AP BC006302

Size: GenelD (NCBI):

150ul , Concentration: 600 ug/ml by 11075

Nanodrop; UNIPROT ID: Source: Q93045

Rabbit Full Name:
Isotype: stathmin-like 2
IgG Calculated MW:

Immunogen Catalog Number: 21 kDa

AG0929 Observed MW:

Observed MV 18-21 kDa

Positive Controls:

Tested Applications: Positive Controls:

WB, IHC, IF/ICC, IP, ELISA

WB: mouse brain tissue, SH-SY5Y cells

Cited Applications:

WB, IHC, IF

Species Specificity:
human. mouse, rat

IP: mouse brain tissue,
IHC: mouse brain tissue,
IHC: mouse brain tissue,
IF/ICC: SH-SY5Y cells,

human, mouse, rat, chicken, zebrafish

human, mouse, rat Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Stathmin-2 (encoded by STMN2), a member of the stathmin family of proteins, encodes a phosphoprotein that has an important role in microtubule dynamics. The STMN2 protein promotes microtubule dynamics in axonal growth cones, and is thought to play an important role in neurite outgrowth. STMN2 protein is localized in both Golgi and cytosolic compartments. Its expression appears modulated in osteoblasts by nerve growth factor, dexamethasone or RhoA kinase inhibitor Y-27632 which are known effectors of osteogenesis. (PMID: 33841129, PMID: 18611392)

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Konstantin Feinberg	28877995	J Cell Biol	WB
Adam Tuttle	31476133	Elife	IHC
Andrea Loreto	31740269	Neurobiol Dis	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 10586-1-AP (STMN2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 10586-1-AP (STMN2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of STMN2 antibody (10586-1-AP; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-STMN2 transfected SH-SY5Y cells.



IP result of anti-STMN2 (IP:10586-1-AP, 4ug; Detection:10586-1-AP 1:10000) with mouse brain tissue lysate 1840 ug.



Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using STMN2 antibody (10586-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).