## For Research Use Only

## CSNK2A2 Polyclonal antibody

Catalog Number: 10606-1-AP 5 Publications



**Purification Method:** 

WB 1:5000-1:50000

IF/ICC 1:200-1:800

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

**Basic Information** 

Catalog Number:

GenBank Accession Number:

BC008812

Size:

GeneID (NCBI):

150ul, Concentration: 500 ug/ml by Nanodrop:

10606-1-AP

**UNIPROT ID:** P19784

Rabbit

Full Name:

Isotype: IgG

casein kinase 2, alpha prime

Immunogen Catalog Number:

polypeptide Calculated MW:

AG0877

41 kDa

Observed MW: 30-35 kDa

**Applications** 

**Tested Applications:** 

WB, IF/ICC, IP, ELISA

**Cited Applications:** WB, IHC, IP

Species Specificity:

human

Cited Species:

human, mouse

Positive Controls:

WB: HEK-293 cells, HepG2 cells, Jurkat cells

IP: Jurkat cells,

IF/ICC: HepG2 cells,

**Background Information** 

CSNK2A2(Casein kinase II subunit alpha') is also named as CK2A3, CK2A2, CSNK2A1, CK2 and belongs to the protein kinase superfamily. It can function as a component of the p53/TP53-dependent spindle assembly checkpoint (SAC) that maintains cyclin-B-CDK1 activity and G2 arrest in response to spindle damage during mitosis. It also can protect against ionizing radiation-induced apoptosis partly by phosphorylating and activating SIRT1. This protein has molecular weights between 35 kDa and 44 kDa(PMID:8622692).

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Dahyun Yu	35659303	Acta Neuropathol Commun	WB,IHC
Yan Zhang	31987502	Biochem Biophys Res Commun	WB
Libin Xu	38909494	Int Immunopharmacol	WB

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 10606-1-AP (CSNK2A2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CSNK2A2 antibody (10606-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



IP result of anti-CSNK2A2 (IP:10606-1-AP, 4ug; Detection:10606-1-AP 1:8000) with Jurkat cells lysate 1400 ug.