

REDD1 specific Polyclonal ANTIBODY

Catalog Number: 10638-1-AP

Featured Product

134 Publications

Basic Information

Catalog Number:
10638-1-AP

Size:
80 µg/150 µl

Source:
Rabbit

Isotype:
IgG

Purification Method:
Antigen affinity purification

Immunogen Catalog Number:
AG0965

GenBank Accession Number:
BC007714

GeneID (NCBI):
54541

Full Name:
DNA-damage-inducible transcript 4

Calculated MW:
25 kDa

Observed MW:
35 kDa

Recommended Dilutions:

WB : 1:500-1:5000

IP : 1:200-1:2000

IHC : 1:50-1:200

Applications

Tested Applications:
ELISA, IP, WB, IHC

Cited Applications:
IF, IHC, IP, WB

Species Specificity:
human, mouse, rat

Cited Species:
human, Meriones unguiculatus, mouse, pig, rabbit, rat

Positive Controls:

WB : K-562 cells; PC-3 cells

IP : MCF-7 cells;

IHC : human kidney tissue; human breast cancer tissue; human gliomas tissue; human heart tissue; human kidney tissue; human liver tissue; human lung cancer tissue; human pancreas cancer tissue; human placenta tissue; human spleen tissue; human testis tissue

Background Information

REDD1, also named as RTP801 and DDIT4, belongs to the DDIT4 family. REDD1 promotes neuronal cell death. It is a novel transcriptional target of p53 implicated ROS in the p53-dependent DNA damage response. REDD1 controlled cell growth under energy stress, as an essential regulator of TOR activity through the TSC1/2 complex. REDD-1 expression has also been linked to apoptosis, Aβ toxicity and the pathogenesis of ischemic diseases. As an HIF-1-responsive gene, REDD-1 exhibits strong hypoxia-dependent upregulation in ischemic cells of neuronal origin [PMD: 19996311]. In response to stress due to DNA damage and glucocorticoid treatment, REDD-1 is upregulated at the transcriptional level [PMD: 21733849]. REDD-1 negatively regulates the mammalian target of Rapamycin, a serine/threonine kinase often referred to as mTOR [PMD: 22951983]. It is crucial in the coupling of extra- and intracellular cues to mTOR regulation. The absence of REDD-1 is associated with the development of retinopathy, a major cause of blindness [PMD: 22304497]. REDD1 is a new host defense factor, and chemical activation of REDD1 expression represents a potent antiviral intervention strategy [PMD: 21909097]. The calculated molecular weight of REDD1 is 25 kDa. Because of multiple lysines in the proteins, REDD1 often migrates around 35 kDa on Western blot [PMD: 19221489]. This antibody is a rabbit polyclonal antibody raised against full length human REDD1 antigen. This antibody is specific to the REDD1 from siRNA experiment (PMD:24713927)

Notable Publications

Author	Pubmed ID	Journal	Application
King Frank W FW	19789631	PLoS One	WB
B Morquette	25257176	Cell Death Differ	WB
Jennifer L Steiner	26394774	Alcohol Alcohol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of REDD1 antibody (10638-1-AP 1:1000) with si-control and si-REDD1 transfected PC-3 cells.



K-562 cells were subjected to SDS PAGE followed by western blot with 10638-1-AP(REDD1 antibody) at dilution of 1:1000



IP Result of anti-REDD1 (IP:10638-1-AP, 3ug; Detection:10638-1-AP 1:500) with MCF-7 cells lysate 2500ug.



Immunohistochemical of paraffin-embedded human kidney using 10638-1-AP(REDD1 antibody) at dilution of 1:100 (under 40x lens)