

For Research Use Only

# NRAS Polyclonal antibody

Catalog Number: 10724-1-AP

Featured Product

33 Publications



## Basic Information

### Catalog Number:

10724-1-AP

### Size:

150ul, Concentration: 273 ug/ml by Bradford method using BSA as the standard;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG1081

### GenBank Accession Number:

BC005219

### GeneID (NCBI):

4893

### UNIPROT ID:

P01111

### Full Name:

neuroblastoma RAS viral (v-ras) oncogene homolog

### Calculated MW:

21 kDa

### Observed MW:

21 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

## Applications

### Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

### Cited Applications:

WB, IHC, IF, IP

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

WB : A431 cells, C6 cells, NIH/3T3 cells, Jurkat cells, A549 cells, HEK-293 cells, MCF-7 cells

IP : HEK-293 cells,

IHC : human pancreas tissue, human testis tissue

IF/ICC : HeLa cells,

## Background Information

NRAS, also named as N-ras and NRAS1, is neuroblastoma RAS viral (v-ras) oncogene homolog from the mammalian ras gene family and it is a member of the small GTPase superfamily. RAS proteins are involved in signal transduction pathways, and bind GDP/GTP and possess intrinsic GTPase activity. It is mapped on chromosome 1, and it is activated in HL60, a promyelocytic leukemia line. Defects in NRAS are a cause of juvenile myelomonocytic leukemia (JMML).

## Notable Publications

Author	Pubmed ID	Journal	Application
Andrew M Waters	28951536	Sci Signal	WB
Akiya Tatsumi	36358254	Biology (Basel)	WB
Chen Liu	33046519	Clin Cancer Res	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10724-1-AP (NRAS antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human pancreas tissue slide using 10724-1-AP (NRAS Antibody) at dilution of 1:200 (under 10x lens).




Immunohistochemical analysis of paraffin-embedded human pancreas tissue slide using 10724-1-AP (NRAS Antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using NRAS antibody (10724-1-AP) at dilution of 1:400 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).



1x10<sup>6</sup> HepG2 cells were intracellularly stained with 0.25 ug NRAS Polyclonal antibody (10724-1-AP) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



IP result of anti-NRAS (IP:10724-1-AP, 4ug; Detection:10724-1-AP 1:4000) with HEK-293 cells lysate 1720 ug.