## For Research Use Only

## Prohibitin Polyclonal antibody

Catalog Number:10787-1-AP

Featured Product

24 Publications




Basic Information	Catalog Number: 10787-1-AP	GenBank Accession Number: BC013401	Purification Metho Antigen affinity p		
	Size:	GenelD (NCBI):	6 71	Recommended Dilutions:	
	150ul , Concentration: 500 ug/ml by Nanodrop; Source:	5245	WB 1:2000-1:1000	WB 1:2000-1:10000 IHC 1:50-1:500 IF/ICC 1:200-1:800	
		UNIPROT ID:			
		P35232	IF/ICC 1:200-1:80		
	Rabbit	Full Name:			
	Isotype: IgG Immunogen Catalog Number: AG1240	prohibitin			
		Calculated MW: 30 kDa			
		Observed MW: 30 kDa			
Applications	Tested Applications:	Positive	Positive Controls:		
	WB, IHC, IF/ICC, ELISA WB :		431 cells, human liver tissue, human brain		
	Cited Applications: WB, IHC, IF, IP, CoIP	AC, IF, IP, ColP cells, mouse brain tissue, mouse hear   es Specificity: kidney tissue, rat brain tissue, rat hear   n, mouse, rat kidney tissue		T3 cells, HeLa cells, Raji cells, Jurkat brain tissue, mouse heart tissue, mous	
	Species Specificity: human, mouse, rat			theart tissue, rat	
	Cited Species:			uman heart tissue	
	human, mouse, rat IF/ICC : HeLa cells, HepG2 cells				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
	bujjer pr 0.0				
Background Information	Prohibitin, also known as PHB, a mer yeast, protozoa and mammals and is ). This protein localizes to the inner r found in the nucleus, where it negati expression and distribution of PHB, w localization with specific gene produ	important in cell proliferation, c nembrane of mitochondria, wher vely regulates transcription (PMI /hich is a nuclear matrix protein,	ifferentiation and apopto e it acts as a chaperone p D: 18558096). Recent stuc affect the apoptosis of Ha	osis (PMID: 2084058 rotein, and is also dy confirmed that th	
	Prohibitin, also known as PHB, a mer yeast, protozoa and mammals and is ). This protein localizes to the inner r found in the nucleus, where it negati expression and distribution of PHB, w localization with specific gene produ	important in cell proliferation, c nembrane of mitochondria, wher vely regulates transcription (PMI /hich is a nuclear matrix protein,	ifferentiation and apopto e it acts as a chaperone p D: 18558096). Recent stuc affect the apoptosis of Ha	osis (PMID: 2084058 rotein, and is also dy confirmed that th	
	Prohibitin, also known as PHB, a mer yeast, protozoa and mammals and is ). This protein localizes to the inner r found in the nucleus, where it negati expression and distribution of PHB, w localization with specific gene produced Author Pub	important in cell proliferation, c nembrane of mitochondria, wher vely regulates transcription (PMI /hich is a nuclear matrix protein, lcts connected with cell apoptosi	ifferentiation and apopto e it acts as a chaperone p D: 18558096). Recent stuc affect the apoptosis of Ha	sis (PMID: 2084058 rotein, and is also dy confirmed that th CaT cells and its co	
Background Information Notable Publications	Prohibitin, also known as PHB, a mer yeast, protozoa and mammals and is ). This protein localizes to the inner r found in the nucleus, where it negati expression and distribution of PHB, w localization with specific gene produ Author Pub Huijing Yin 315	important in cell proliferation, c nembrane of mitochondria, wher vely regulates transcription (PMI /hich is a nuclear matrix protein, icts connected with cell apoptosi med ID Journal 01431 Cell Death Dis	ifferentiation and apopto e it acts as a chaperone p D: 18558096). Recent stuc affect the apoptosis of Ha	sis (PMID: 2084058 rotein, and is also dy confirmed that th CaT cells and its co Application	
	Prohibitin, also known as PHB, a mer yeast, protozoa and mammals and is ). This protein localizes to the inner r found in the nucleus, where it negati expression and distribution of PHB, v localization with specific gene produ Author Pub Huijing Yin 315 Chong-Chong Guo 265	important in cell proliferation, c nembrane of mitochondria, wher vely regulates transcription (PMI /hich is a nuclear matrix protein, icts connected with cell apoptosi med ID Journal 01431 Cell Death Dis	ifferentiation and apopto e it acts as a chaperone pi D: 18558096). Recent stuc affect the apoptosis of Ha s (PMID: 24402549).	sis (PMID: 2084058 rotein, and is also dy confirmed that th CaT cells and its co Application WB	
	Prohibitin, also known as PHB, a mer yeast, protozoa and mammals and is ). This protein localizes to the inner r found in the nucleus, where it negati expression and distribution of PHB, v localization with specific gene produ Author Pub Huijing Yin 315 Chong-Chong Guo 265	important in cell proliferation, c nembrane of mitochondria, wher vely regulates transcription (PMI /hich is a nuclear matrix protein, icts connected with cell apoptosi med ID Journal 01431 Cell Death Dis 57149 Evid Based Corr 45295 EMBO J	ifferentiation and apopto e it acts as a chaperone pi D: 18558096). Recent stuc affect the apoptosis of Ha s (PMID: 24402549).	sis (PMID: 2084058 rotein, and is also dy confirmed that th CaT cells and its co Application WB WB	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10787-1-AP (Prohibitin antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 10787-1-AP (Prohibitin antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 10787-1-AP (Prohibitin antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 10787-1-AP (Prohibitin antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Prohibitin antibody (10787-1-AP) at dilution of 1:400 and Multi-rAb Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).