For Research Use Only

Cyclin D3 Polyclonal antibody

Catalog Number: 10845-1-AP 6 Publications



Basic Information	Catalog Number: 10845-1-AP	GenBank Accession Number: BC011616	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Anagenanning partication
	150ul , Concentration: 133 µg/ml by	896	
	Bradford method using BSA as the	ENSEMBL Gene ID:	
	standard;	ENSG00000112576	
	Source: Rabbit	UNIPROT ID: P30281	
	Isotype: IgG	Full Name: cyclin D3	
	Immunogen Catalog Number: AG1292	Calculated MW: 33 kDa	
Applications	Tested Applications: WB, IF, IHC, ELISA		
	Cited Applications: WB, IHC		
	Species Specificity: human		
	Cited Species:		
Background Information	human Cyclin D3 belongs to the highly cons	erved cyclin family, whose memb	ers are characterized by a dramatic
Background Information	Cyclin D3 belongs to the highly cons periodicity in protein abundance thro cyclins exhibit distinct expression ar mitotic event. This cyclin forms a cor activtiy is required for cell cycle G1/ phosphorylation of tumor suppressor	bugh the cell cycle. Cyclins function nd degradation patterns which con mplex with and functions as a regu S transition. This protein has been protein Rb. The CDK4 activity ass	n as regulators of CDK kinases. Different tribute to the temporal coordination of each ılatory subunit of CDK4 or CDK6, whose shown to interact with and be involved in th ociated with this cyclin was reported to be
Background Information	Cyclin D3 belongs to the highly cons periodicity in protein abundance thro cyclins exhibit distinct expression ar mitotic event. This cyclin forms a cor activtiy is required for cell cycle G1/2	bugh the cell cycle. Cyclins function nd degradation patterns which con mplex with and functions as a regu S transition. This protein has been protein Rb. The CDK4 activity ass	n as regulators of CDK kinases. Different tribute to the temporal coordination of each ılatory subunit of CDK4 or CDK6, whose shown to interact with and be involved in th ociated with this cyclin was reported to be
	Cyclin D3 belongs to the highly cons periodicity in protein abundance thro cyclins exhibit distinct expression ar mitotic event. This cyclin forms a cor activity is required for cell cycle G1/2 phosphorylation of tumor suppressor necessary for cell cycle progression t	bugh the cell cycle. Cyclins function nd degradation patterns which con mplex with and functions as a regu S transition. This protein has been protein Rb. The CDK4 activity ass	n as regulators of CDK kinases. Different tribute to the temporal coordination of each ılatory subunit of CDK4 or CDK6, whose shown to interact with and be involved in th ociated with this cyclin was reported to be
	Cyclin D3 belongs to the highly cons periodicity in protein abundance thro cyclins exhibit distinct expression ar mitotic event. This cyclin forms a cor activity is required for cell cycle G1/ phosphorylation of tumor suppressor necessary for cell cycle progression t	bugh the cell cycle. Cyclins function nd degradation patterns which con mplex with and functions as a regu S transition. This protein has been protein Rb. The CDK4 activity asso hrough G2 phase into mitosis afte	n as regulators of CDK kinases. Different tribute to the temporal coordination of each Ilatory subunit of CDK4 or CDK6, whose shown to interact with and be involved in th ociated with this cyclin was reported to be r UV radiation.
	Cyclin D3 belongs to the highly cons periodicity in protein abundance thro cyclins exhibit distinct expression ar mitotic event. This cyclin forms a cor activity is required for cell cycle G1/2 phosphorylation of tumor suppressor necessary for cell cycle progression t Author Pub Zhang Ping P 169	bugh the cell cycle. Cyclins function and degradation patterns which com mplex with and functions as a regu S transition. This protein has been protein Rb. The CDK4 activity asso hrough G2 phase into mitosis after med ID Journal	n as regulators of CDK kinases. Different tribute to the temporal coordination of each llatory subunit of CDK4 or CDK6, whose shown to interact with and be involved in th ociated with this cyclin was reported to be r UV radiation. Application
	Cyclin D3 belongs to the highly consperiodicity in protein abundance thro cyclins exhibit distinct expression ar mitotic event. This cyclin forms a cor activity is required for cell cycle G1/2 phosphorylation of tumor suppressor necessary for cell cycle progression to Author Pub Zhang Ping P 169 Wenkai Xiao 253	bugh the cell cycle. Cyclins function ad degradation patterns which con mplex with and functions as a reguly S transition. This protein has been protein Rb. The CDK4 activity association hrough G2 phase into mitosis after med ID Journal 78399 BMC Cancer	n as regulators of CDK kinases. Different tribute to the temporal coordination of each llatory subunit of CDK4 or CDK6, whose shown to interact with and be involved in th ociated with this cyclin was reported to be r UV radiation. Application
Notable Publications	Cyclin D3 belongs to the highly consperiodicity in protein abundance thro cyclins exhibit distinct expression ar mitotic event. This cyclin forms a cor activity is required for cell cycle G1/2 phosphorylation of tumor suppressor necessary for cell cycle progression to Author Pub Zhang Ping P 169 Wenkai Xiao 253 Mingmin Tang 330	bugh the cell cycle. Cyclins function nd degradation patterns which con mplex with and functions as a regul S transition. This protein has been protein Rb. The CDK4 activity assistion hrough G2 phase into mitosis after med ID Journal 78399 BMC Cancer 55542 Int J Oncol 191855 Phytomedicine	n as regulators of CDK kinases. Different tribute to the temporal coordination of each llatory subunit of CDK4 or CDK6, whose shown to interact with and be involved in th ociated with this cyclin was reported to be r UV radiation. Application WB,IHC
Notable Publications	Cyclin D3 belongs to the highly consi periodicity in protein abundance thro cyclins exhibit distinct expression ar mitotic event. This cyclin forms a cor activity is required for cell cycle G1/2 phosphorylation of tumor suppressor necessary for cell cycle progression t Author Pub Zhang Ping P 169 Wenkai Xiao 253 Mingmin Tang 330 Storage: Stora at -20°C. Stable for one year afte Storage Buffer:	bugh the cell cycle. Cyclins function nd degradation patterns which con mplex with and functions as a reguly S transition. This protein has been protein Rb. The CDK4 activity assistion hrough G2 phase into mitosis after med ID Journal 78399 BMC Cancer 55542 Int J Oncol 191855 Phytomedicine	n as regulators of CDK kinases. Different tribute to the temporal coordination of each llatory subunit of CDK4 or CDK6, whose shown to interact with and be involved in th ociated with this cyclin was reported to be r UV radiation. Application WB,IHC
Notable Publications	Cyclin D3 belongs to the highly consperiodicity in protein abundance thro cyclins exhibit distinct expression ar mitotic event. This cyclin forms a cor activity is required for cell cycle G1/2 phosphorylation of tumor suppressor necessary for cell cycle progression t Author Pub Zhang Ping P 169 Wenkai Xiao 253 Mingmin Tang 330 Storage: Store at -20°C. Stable for one year after	bugh the cell cycle. Cyclins function nd degradation patterns which con nplex with and functions as a reguly S transition. This protein has been protein Rb. The CDK4 activity assist hrough G2 phase into mitosis after med ID Journal 78399 BMC Cancer 55542 Int J Oncol 191855 Phytomedicine ter shipment. P% glycerol pH 7.3.	n as regulators of CDK kinases. Different tribute to the temporal coordination of each llatory subunit of CDK4 or CDK6, whose shown to interact with and be involved in th ociated with this cyclin was reported to be r UV radiation. Application WB,IHC
Notable Publications	Cyclin D3 belongs to the highly consperiodicity in protein abundance thro cyclins exhibit distinct expression ar mitotic event. This cyclin forms a cor activity is required for cell cycle G1/2 phosphorylation of tumor suppressor necessary for cell cycle progression t Author Pub Zhang Ping P 169 Wenkai Xiao 253 Mingmin Tang 330 Storage: Stora et -20°C. Stable for one year aff Storage Buffer: PBS with 0.02% sodium azide and 50	bugh the cell cycle. Cyclins function nd degradation patterns which con nplex with and functions as a reguly S transition. This protein has been protein Rb. The CDK4 activity assist hrough G2 phase into mitosis after med ID Journal 78399 BMC Cancer 55542 Int J Oncol 191855 Phytomedicine ter shipment. P% glycerol pH 7.3.	n as regulators of CDK kinases. Different tribute to the temporal coordination of each llatory subunit of CDK4 or CDK6, whose shown to interact with and be involved in th ociated with this cyclin was reported to be r UV radiation. Application WB,IHC

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer. Selected Validation Data