

For Research Use Only

# STK38 Polyclonal antibody

Catalog Number: 11105-1-AP

Featured Product

1 Publications



## Basic Information

### Catalog Number:

11105-1-AP

### Size:

150ul, Concentration: 450 ug/ml by Nanodrop;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG1559

### GenBank Accession Number:

BC012085

### GeneID (NCBI):

11329

### UNIPROT ID:

Q15208

### Full Name:

serine/threonine kinase 38

### Calculated MW:

54 kDa

### Observed MW:

50-55 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

## Applications

### Tested Applications:

WB, IHC, IP, ELISA

### Cited Applications:

WB, IP

### Species Specificity:

human, mouse

### Cited Species:

human

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB:** HEK-293 cells, HeLa cells, HepG2 cells, Jurkat cells, MCF-7 cells

**IP:** MCF-7 cells,

**IHC:** mouse cerebellum tissue,

## Background Information

Serine/threonine-protein kinase 38 belongs to AGC Ser/Thr protein kinase family. STK38 implicates in cell proliferation and/or tumor progression, and might play a role in the HIV-1 life cycle. Researches showed that STK38 with STK38L, were proapoptotic kinases and key members of the RASSF1/STK4 signaling cascade and its kinase activity increased in early mitotic phase and was dependent on FRY and STK3. This antibody recognizes the C-terminal of STK38.

## Notable Publications

Author	Pubmed ID	Journal	Application
Xin Li	28361952	Nat Commun	WB, IP

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of STK38 antibody (11105-1-AP; 1:400; incubated at room temperature for 1.5 hours) with sh-Control and sh-STK38 transfected HeLa cells.




IP result of anti-STK38 (IP:11105-1-AP, 4ug; Detection:11105-1-AP 1:300) with MCF-7 cells lysate 800ug.



Various lysates were subjected to SDS PAGE followed by western blot with 11105-1-AP (STK38 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue slide using 11105-1-AP (STK38 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue slide using 11105-1-AP (STK38 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).