For Research Use Only

TCEAL7 Polyclonal antibody

Catalog Number:11218-1-AP

Featured Product

2 Publications



Basic Information

Catalog Number: GenBank Accession Number: 11218-1-AP

Antigen affinity purification BC016786 GeneID (NCBI): Recommended Dilutions: Size

WB 1:500-1:1000 150ul, Concentration: 700 µg/ml by 56849 Nanodrop and 300 µg/ml by Bradford Full Name: IHC 1:20-1:200 method using BSA as the standard;

transcription elongation factor A (SII)-

Rabbit

Calculated MW: 12 kDa Isotype: IgG Observed MW: Immunogen Catalog Number: 12 kDa

AG1715

Applications

Tested Applications: IHC, WB, ELISA

Cited Applications:

IHC, WB

Species Specificity:

human **Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

WB: human liver tissue, A549 cells

Positive Controls:

IHC: human ovary tumor tissue, human gliomas tissue

Purification Method:

Background Information

TCEAL7 is a transcription regulatory factor that has a role in the negative regulation of NF-kappa-B signaling at the basal level by regulating transcriptional activity of NF-kappa-B on its target gene promoters. It can associate with cyclin D1 promoter containing Myc E-box sequence and transcriptionally represses cyclin D1 expression. In addition, TCEAL7 regulates telomerase reverse transcriptase expression and telomerase activity in both ALT (alternative lengthening of telomeres) and telomerase-positive cell lines

Notable Publications

Author	Pubmed ID	Journal	Application
Huang Chun-yu CY	23372750	PLoS One	WB,IHC
Xilin Liu	33430878	Cancer Cell Int	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



human liver tissue were subjected to SDS PAGE followed by western blot with 11218-1-AP (TCEAL7 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary tumor using 11218-1-AP (TCEAL7 antibody) at dilution of 1:50 (under 40x lens).