For Research Use Only

RARG Polyclonal antibody

Catalog Number: 11424-1-AP

Featured Product

6 Publications



Basic Information

Catalog Number: GenBank Accession Number:

 11424-1-AP
 BC019098

 Size:
 GeneID (NCBI):

150ul , Concentration: 1000 ug/ml by 5916
Nanodrop and 467 ug/ml by Bradford
method using BSA as the standard;
Source: UNIPROT ID:
p13631
Source: Full Name of

Source: Full Name:
Rabbit retinoic acid receptor, gamma

Isotype: Calculated MW:
IgG 454 aa, 50 kDa
Immunogen Catalog Number: Observed MW:
AG1991 48-50 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

Applications

Tested Applications:

WB, IP, ELISA

Cited Species: human, mouse

Cited Applications: WB, IHC, IF, CoIP, ChIP Species Specificity: human, mouse, rat Positive Controls:

WB: HeLa cells, MCF-7 cells

IP: MCF-7 cells,

Background Information

Retinoic acid receptors (RARs) are nuclear hormone receptors that act as ligand-dependent transcriptional regulators. In their liganded state, RARs activate transcription, whereas in their nonliganded form, they repress transcription of their target genes. RARs have numerous target genes, which have retinoic response elements in their promoter regions [PMID:17574023]. RARG is a member of the retinoid acid receptor (RAR) family, along with RARA, which is known to be involved PML/RARA fusion of acute promyelocytic leukemia, and RARB. and rearrangement that conferred to leukemic cells acute promyelocytic leukemia-like morphologic and immunophenotypic features [PMID:20935257].

Notable Publications

Author	Pubmed ID	Journal	Application
Lin Xiu	36523991	Front Oncol	WB,IHC
Tomoko Kawai	27022678	Endocrinology	IF
Yao Zhang	36031407	Adv Sci (Weinh)	ChIP

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 11424-1-AP (RARG antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



IP result of anti-RARG (IP:11424-1-AP, 4ug; Detection:11424-1-AP 1:300) with MCF-7 cells lysate 1120ug.