For Research Use Only

SESN3 Polyclonal antibody

Catalog Number: 11431-2-AP

Featured Product

18 Publications



Basic Information

Catalog Number:

11431-2-AP

Size:

150ul, Concentration: 173 ug/ml by 143686 Bradford method using BSA as the

standard;

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG1961

GenBank Accession Number:

BC017296 GeneID (NCBI):

UNIPROT ID:

P58005 Full Name: sestrin 3

Calculated MW: 492 aa, 57 kDa Observed MW:

36 kDa, 51-59 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, IHC

Species Specificity: human, mouse, rat **Cited Species:** human, mouse, rat

Positive Controls:

WB: HeLa cells, A431 cells, A549 cells, HEK-293 cells, HepG2 cells, mouse liver tissue, mouse ovary tissue

Background Information

Sestrins, including sestrin-1 (PA26), sestrin-2 (Hi95), and sestrin-3, are 48 to 65 kDa cystein sulfinyl reductases and they modulate peroxide signaling and antioxidant defense. These proteins selectively reduce or repair hyperoxidized forms of typical 2-Cys peroxiredoxins within eukaryotes. Expression of these proteins is regulated by p53, a tumor suppressor protein. Sestrin 3 was identified as a forkhead box O (FoxO) target gene with antioxidant activity. Recently it was reported that sestrin 3 may play an important role in AKT induced increase in ROS and it might be a promising target in selectively killing cancer cells containing high levels of AKT activity.

Notable Publications

Author	Pubmed ID	Journal	Application
T Shen	27693501	Osteoarthritis Cartilage	WB,IHC
Li-Gang Huang	29191771	Brain Res	WB
Vakana Eliza E	24260131	PLoS One	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 11431-2-AP (SESN3 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.