### For Research Use Only

# COX4I2 Polyclonal antibody

Catalog Number: 11463-1-AP

Featured Product

19 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

 11463-1-AP
 BC057779

 Size:
 GeneID (NCBI):

 150ul , Concentration: 850 ug/ml by
 84701

150ul , Concentration: 850 ug/ml by 84701
Nanodrop and 333 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; 006K10

method using BSA as the standard; Q96KJ9

Source: Full Name:

Rabbit cytochrome c oxidase subunit IV Isotype: isoform 2 (lung)

IgG Calculated MW:
Immunogen Catalog Number: 171 aa, 20 kDa
AG1987 Observed MW:

17-18 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB: 1:500-1:3000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC: 1:20-1:200 IF/ICC: 1:200-1:800

**Applications** 

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

**Cited Applications:** 

WB, IHC, IF

Species Specificity: human, mouse, rat Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HepG2 cells, human brain tissue, MCF-7 cells, human heart tissue, human lung tissue, A375 cells

IP: HepG2 cells,

IHC: human ovary tumor tissue, IF/ICC: HepG2 cells, HeLa cells

## Background Information

COX4I2, also named as COX4L2 and COXIV-2, belongs to the cytochrome c oxidase IV family. It is one of the nuclear-coded polypeptide chains of cytochrome c oxidase, the terminal oxidase in mitochondrial electron transport. COXIV(cytochrome c oxidase IV) has two isoforms (isoform 1 and 2). Isoform 1(COX4I1) is ubiquitously expressed and isoform 2 is highly expressed in lung tissues. This antibody was generated against full length COX4I2 protein.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yongxin Mao	36172142	Front Oncol	IHC,IF
Siddhesh Aras	27913209	Biochim Biophys Acta	WB
Guy Keller	30423473	Neurobiol Dis	IF

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (foll free F: proteintech@ptglab.com


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


### Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11463-1-AP (COX412 antibody) at dilution of 1:1500 incubated at 4 degree celsius over night.




IP result of anti-COX4I2 (11463-1-AP for IP and Detection) with HepG2 cell lysate.



IP result of anti-COX4I2 (IP:11463-1-AP, 3ug; Detection:11463-1-AP 1:1000) with HepG2 cells lysate 8000ug.



Immunohistochemical analysis of paraffinembedded human ovary tumor using 11463-1-AP (COX412 antibody) at dilution of 1:50 (under 10x lens).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using COX412 antibody (11463-1-AP) at dilution of 1:400 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using COX4/2 antibody (11463-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).