

For Research Use Only

FXYP7 Polyclonal antibody

Catalog Number:11465-1-AP



Basic Information

Catalog Number:

11465-1-AP

Size:

150ul , Concentration: 650 ug/ml by Nanodrop and 320 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG2030

GenBank Accession Number:

BC018619

GeneID (NCBI):

53822

UNIPROT ID:

P58549

Full Name:

FXYP domain containing ion transport regulator 7

Calculated MW:

9 kDa

Observed MW:

18-20 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human, mouse, rat, pig

Positive Controls:

WB : mouse brain tissue, human brain tissue, rat brain tissue

Background Information

FXYP7 (FXYP domain containing ion transport regulator 7) is a single-pass membrane protein that belongs to the FXYP family. The FXYP family, which contains seven members, are tissue specific regulators of the Na,K-ATPase. Expressed exclusively in the brain, FXYP7 is an isoform-specific Na,K-ATPase regulator which could play an important role in neuronal excitability. It has been reported that FXYP7 bears post-translationally added modifications on threonine residues and has an apparent molecular weight of 18 kDa (PMID: 12093728).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 11465-1-AP (FXVD7 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.