

For Research Use Only

# OGT Polyclonal antibody

Catalog Number: 11576-2-AP

Featured Product

68 Publications



## Basic Information

### Catalog Number:

11576-2-AP

### Size:

150ul, Concentration: 800 ug/ml by Nanodrop and 367 ug/ml by Bradford method using BSA as the standard;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG2160

### GenBank Accession Number:

BC014434

### GeneID (NCBI):

8473

### UNIPROT ID:

O15294

### Full Name:

O-linked N-acetylglucosamine (GlcNAc) transferase (UDP-N-acetylglucosamine:polypeptide-N-acetylglucosaminyl transferase)

### Calculated MW:

1046 aa, 117 kDa

### Observed MW:

110 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:2000-1:12000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

## Applications

### Tested Applications:

WB, IHC, IP, ELISA

### Cited Applications:

WB, IHC, IF, IP, CoIP, IP-MS

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse, rat

### Positive Controls:

**WB**: HepG2 cells, mouse liver tissue, mouse brain tissue, rat brain tissue

**IP**: mouse brain tissue,

**IHC**: human colon cancer tissue, human lung cancer tissue, human pancreas cancer tissue, rat testis tissue

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

O-linked N-acetylglucosamine transferase (OGT) catalyzes the attachment of N-acetylglucosamine (GlcNAc) monosaccharides to the hydroxyl group of serine or threonine residues of numerous nuclear and cytoplasmic proteins and may play important roles in a large number of diverse intracellular processes ranging from translational control, transcription, transcriptional repression, INS resistance and regulation of the cell cycle. It exists as a heterotrimeric complex with two 110 kDa and one 70 kDa subunits. Recent studies have shown that O-GlcNAcylation plays essential roles in cancer formation and progression. O-GlcNAcylation as well as OGT expression was found to be significantly elevated in the cancer tissues.

## Notable Publications

Author	Pubmed ID	Journal	Application
Xiao Han	31545463	Oncol Rep	
Jing Zhang	31539718	Atherosclerosis	WB
Chia-Wei Hu	29058723	Nat Chem Biol	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 11576-2-AP (OGT antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 11576-2-AP (OGT antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



IP result of anti-OGT (IP:11576-2-AP, 3ug; Detection:11576-2-AP 1:1000) with mouse brain tissue lysate 8000ug.



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 11576-2-AP (OGT antibody) at dilution of 1:300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).