For Research Use Only

## LY96/MD2 Polyclonal antibody

Catalog Number:11784-1-AP 13 Publications





Basic Information	Catalog Number: 11784-1-AP	GenBank Accession Number: BC020690	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 550 ug/ml by	23643	WB 1:500-1:2000	
	Nanodrop;	UNIPROT ID:	IHC 1:50-1:500	
	Source: Rabbit	Q9Y6Y9		
		Full Name:		
	Isotype: IgG	lymphocyte antigen 96		
	Immunogen Catalog Number:	Calculated MW: 160 aa, 18 kDa Observed MW: 20-26 kDa		
	AG2447			
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, ELISA WB : mouse		use testis tissue, rat testis tissue	
	Cited Applications: WB, IHC, IF, IP, CoIP	IHC : mouse testis tissue,		
	Species Specificity: human, mouse, rat			
	Cited Species: human, mouse, rat			
	Note-IHC: suggested antigen ( TE buffer pH 9.0; (*) Alternati retrieval may be performed w	vely, antigen		
	buffer pH 6.0			
Background Information	LY96, also named as ESOP1 or MD 2, LPS and to the extracellular domain and induces the dimerization of LPS-	of TLR4. The interaction between MD-2-TLR4, which forms the struc n mediating the interaction betw	that can bind to both the hydrophobic portion of MD-2 and LPS bridges the two TLR4 molecules tural basis for biological functions of TLR4/ME een LPS and TLR4, MD-2 has been extensively lers.	
	LY96, also named as ESOP1 or MD 2, LPS and to the extracellular domain and induces the dimerization of LPS- 2 complex. Due to its essential role i explored as a therapeutic target for t	of TLR4. The interaction between MD-2-TLR4, which forms the struc n mediating the interaction betw	MD-2 and LPS bridges the two TLR4 molecules tural basis for biological functions of TLR4/ME een LPS and TLR4, MD-2 has been extensively	
	LY96, also named as ESOP1 or MD 2, LPS and to the extracellular domain and induces the dimerization of LPS- 2 complex. Due to its essential role i explored as a therapeutic target for t Author Put	of TLR4. The interaction between MD-2-TLR4, which forms the struc n mediating the interaction betw reatment of inflammatory disord	MD-2 and LPS bridges the two TLR4 molecules tural basis for biological functions of TLR4/MD een LPS and TLR4, MD-2 has been extensively lers. Application	
Background Information	LY96, also named as ESOP1 or MD 2, LPS and to the extracellular domain and induces the dimerization of LPS- 2 complex. Due to its essential role i explored as a therapeutic target for t Author Put Min Zhang 364	of TLR4. The interaction between MD-2-TLR4, which forms the struc n mediating the interaction betw reatment of inflammatory disord med ID Journal	MD-2 and LPS bridges the two TLR4 molecules tural basis for biological functions of TLR4/MD een LPS and TLR4, MD-2 has been extensively lers. Application	
	LY96, also named as ESOP1 or MD 2, LPS and to the extracellular domain and induces the dimerization of LPS- 2 complex. Due to its essential role i explored as a therapeutic target for t Author Put Min Zhang 364 Mengya Zhao 353	of TLR4. The interaction between MD-2-TLR4, which forms the struc n mediating the interaction betw reatment of inflammatory disord med ID Journal s14003 J Agric Food Ch	MD-2 and LPS bridges the two TLR4 molecules tural basis for biological functions of TLR4/ML een LPS and TLR4, MD-2 has been extensively lers.	
	LY96, also named as ESOP1 or MD 2, LPS and to the extracellular domain and induces the dimerization of LPS- 2 complex. Due to its essential role i explored as a therapeutic target for t Author Put Min Zhang 364 Mengya Zhao 353	of TLR4. The interaction between MD-2-TLR4, which forms the struc n mediating the interaction betw reatment of inflammatory disord med ID Journal (14003 J Agric Food Ch (172062 Front Oncol (157279 J Cardiovasc Tr (1976) Structure of the structure of the structure (1976) Structure of the structure of the structure of the structure (1976) Structure of the structure of the structure of the structure (1976) Structure of the structu	MD-2 and LPS bridges the two TLR4 molecules tural basis for biological functions of TLR4/ML een LPS and TLR4, MD-2 has been extensively lers.	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 11784-1-AP (LY96/MD2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 11784-1-AP (LY96/MD2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).