

For Research Use Only

Ficolin 3 Polyclonal antibody

Catalog Number: 11867-1-AP

4 Publications



Basic Information

Catalog Number:

11867-1-AP

Size:

150ul, Concentration: 200 ug/ml by Nanodrop and 167 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG2449

GenBank Accession Number:

BC020731

GeneID (NCBI):

8547

UNIPROT ID:

O75636

Full Name:

ficolin (collagen/fibrinogen domain containing) 3 (Hakata antigen)

Calculated MW:

33 kDa

Observed MW:

32 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IF/ICC, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human

Cited Species:

human, mouse, rat

Positive Controls:

WB : human plasma,

IF/ICC : HepG2 cells,

Background Information

Ficolins are a group of oligomeric lectins with subunits consisting of both collagen-like and fibrinogen-like domains (PMID: 20375620). They are innate pattern recognition receptors and play integral roles within the innate immune response. In humans, there are three ficolins termed M-, L-, and H-ficolin (also referred to as ficolin-1, -2, and -3) whereas rodents only possess ficolin-A and -B, which are the orthologues of human L- and M-ficolin, respectively (PMID: 30868077). Ficolin-3 (Hakata antigen or H-ficolin) was initially identified based on its reactivity with sera from patients with systemic lupus erythematosus (PMID: 1859827). It has been shown to have a calcium-independent lectin activity. Ficolin-3 is associated with MASPs and sMAP, and which activates the lectin pathway (PMID: 11907111). Ficolin-3 shows affinity for GalNAc, GlcNAc, D-fucose and galactose (PMID: 30868077).

Notable Publications

Author	Pubmed ID	Journal	Application
Huafeng Song	35300115	Front Mol Biosci	WB
Nadia Houerbi	38862464	Nat Commun	WB
Yanmei Yuan	38698462	J Exp Clin Cancer Res	WB, IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human plasma were subjected to SDS PAGE followed by western blot with 11867-1-AP (Ficolin-3 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Ficolin 3 antibody (11867-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).